

Clinical Policy: Invasive and Noninvasive Home Ventilators

Reference Number: LA.CP.MP.184c

Date of Last Revision: 07/23

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

This policy describes medical necessity criteria for noninvasive and invasive home ventilators. Noninvasive ventilation (NIV) describes the administration of positive pressure to the lungs using interfaces such as, but not limited to, nasal masks, orofacial masks, full face masks, mouthpieces, nasal pillows, or helmets.^{1,2} Invasive ventilatory support describes the administration of positive pressure to the lungs through an invasive interface, such as a tracheostomy tube or endotracheal tube.¹

Policy/Criteria

- I. It is the policy of Louisiana Healthcare Connections that non-invasive *home ventilators* are **medically necessary** for the following indications:
 - A. Initial request for the first three months of noninvasive home ventilator use for restrictive thoracic disorders, **ALL** of the following: (parts 1-4)
 1. Documentation of a neuromuscular disease (ex. amyotrophic lateral sclerosis) or a severe thoracic cage abnormality (ex. post-thoracoplasty for tuberculosis or severe kyphoscoliosis) and both of the following:
 - a. One of the following:
 - i. An arterial blood gas partial pressure of carbon dioxide (PaCO₂) was measured while awake and breathing room air or on prescribed oxygen with a measurement of: PaCO₂ >45 mm Hg;
 - ii. Sleep oximetry demonstrates O₂ saturation of one of the following for at least five minutes while breathing prescribed O₂:
 - a. ≤88% for members/enrollees ≥ 18 years of age;
 - b. <92% for members/enrollees < 18 years of age;
 - b. If neuromuscular disease is present, one of the following:
 - i. For those ≥ 18 years of age, maximal inspiratory pressure is < 60 cm H₂O, or forced vital capacity is < 50% predicted;
 - ii. For those < 18 years of age, documentation of Type 1 (hypoxemic) and/or Type 2 (hypercapnic) respiratory failure or inability to maintain airflow.
 2. Respiratory failure has failed to improve with an adequate trial of bilevel positive airway pressure (Bi-PAP), as evidenced by one of the following: (Note: PaCO₂ levels may not normalize even with adequate response to Bi-PAP therapy. Failure to normalize PaCO₂ levels alone is not considered a therapeutic failure of Bi-PAP)
 - a. Intolerance to Bi-PAP, as indicated by the member's/enrollee's request to discontinue nocturnal assisted ventilation;
 - b. Worsening dyspnea, hemodynamic instability, or unresponsive hypoxemia;
 - c. Signs of respiratory failure, including tachypnea, increased work of breathing, hypoxemia, hypercapnia and/or respiratory acidosis (e.g., pH <7.35);
 3. Chronic obstructive pulmonary disease (COPD) does not contribute significantly to the pulmonary limitation;

4. None of the following contraindications:
 - a. Baseline FIO₂ requirement > 0.40;
 - b. Positive-end expiratory pressure (PEEP) > 10 cm H₂O;
 - c. Need for continuous invasive monitoring in adult patients.

- B. Initial request for the first three months of noninvasive home ventilator use for severe COPD, all of the following: (parts 1-4)
 1. An arterial blood gas PaCO₂ measurement was done while awake and breathing at baseline and prescribed FIO₂, which is greater than or equal to 52 mm Hg;
 2. Prior to initiating therapy, sleep apnea and treatment with a continuous positive airway pressure device (CPAP) has been considered and ruled out. (Note: Formal sleep testing is not required if the medical record demonstrates that sleep apnea (Obstructive Sleep Apnea (OSA), CSA and/or CompSA) is not the predominant cause of awake hypercapnia or nocturnal arterial oxygen desaturation);
 3. Respiratory failure has failed to improve with an adequate trial of Bi-PAP, as evidenced by one of the following: (Note: PaCO₂ levels may not normalize even with adequate response to Bi-PAP therapy. Failure to normalize PaCO₂ levels alone is not considered a therapeutic failure of Bi-PAP);
 - a. Intolerance to Bi-PAP, as indicated by the member's/enrollee's request to discontinue nocturnal assisted ventilation;
 - b. Worsening dyspnea, hemodynamic instability, or unresponsive hypoxemia;
 - c. Signs of respiratory failure, including tachypnea, increased work of breathing, hypoxemia, hypercapnia and/or respiratory acidosis (e.g., pH <7.35);
 4. None of the following contraindications:
 - a. Baseline FIO₂ requirement > 0.40;
 - b. PEEP > 10 cm H₂O;
 - c. Need for continuous invasive monitoring.

- C. Initial request for the first three months of noninvasive home ventilator use for obesity hypoventilation syndrome (OHS) (also known as Pickwickian syndrome), all of the following (parts 1-4):
 1. BMI greater than 30;
 2. An initial arterial blood gas PaCO₂, done while awake and breathing the beneficiary's prescribed FIO₂, is ≥ 45 mm Hg;
 3. Signs of respiratory failure have failed to improve with an adequate trial of Bi-PAP as evidenced by one of the following: (Note: PaCO₂ levels may not normalize even with adequate response to Bi-PAP therapy. Failure to normalize PaCO₂ levels alone is not considered a therapeutic failure of Bi-PAP);
 - a. Intolerance to Bi-PAP, as indicated by the member's/enrollee's request to discontinue nocturnal assisted ventilation;
 - b. Worsening dyspnea, hemodynamic instability, or unresponsive hypoxemia;
 - c. Signs of respiratory failure, including tachypnea, increased work of breathing, hypoxemia, hypercapnia, and/or respiratory acidosis (e.g., pH <7.35).
 - d. An arterial blood gas PaCO₂, done during sleep or immediately upon awakening, and breathing the beneficiary's prescribed FIO₂, shows the member/enrollees

PaCO₂ worsened greater than or equal to 7 mm Hg compared to the original result (see C.2);

4. None of the following contraindications:
 - a. Baseline FIO₂ requirement > 0.40;
 - b. PEEP > 10 cm H₂O;
 - c. Need for continuous invasive monitoring.

D. Initial request for the first three months of noninvasive home ventilator use for members/enrollees who have experienced treatment failure with Bi-PAP, both of the following:

1. Treatment failure, one of the following:
 - a. Intolerance to Bi-PAP, as indicated by member/enrollee request to discontinue nocturnal assisted ventilation;
 - b. Worsening dyspnea, hemodynamic instability, or unresponsive hypoxemia;
 - c. Signs of respiratory failure including tachypnea, increased work of breathing, hypoxemia, hypercapnia and/or respiratory acidosis (e.g., pH <7.35); (PaCO₂ levels may not normalize even with adequate response to Bi-PAP therapy. Failure to normalize PaCO₂ levels alone is not considered a therapeutic failure of Bi-PAP);
2. None of the following contraindications:
 - a. Baseline FIO₂ requirement > 0.40;
 - b. PEEP > 10 cm H₂O;
 - c. Need for continuous invasive monitoring.

II. It is the policy of **Louisiana Healthcare Connections** that *continued use of noninvasive home ventilators* after the initial three months certification period—A Twelve-month certification period is **medically necessary** when meeting the following: (parts A-C)

- A. Medical records document improvement in relevant signs or symptoms due to the device;
- B. The device is used for at least an average of four hours per 24-hour period;
- C. None of the following contraindications:
 1. Baseline FIO₂ requirement > 0.40;
 2. PEEP > 10 cm H₂O;
 3. Need for continuous invasive monitoring.

III. It is the policy Louisiana Healthcare Connections of *that initial and ongoing use of an invasive ventilator* is **medically necessary** for a long-term/chronic condition or disease affecting the ability to effectively maintain an adequate respiratory status. Examples of conditions may include neuromuscular disease, thoracic restrictive disease, or chronic respiratory failure following COPD.

IV. It is the policy of Louisiana Healthcare Connections that *a second or back up noninvasive or invasive ventilator* is considered **medically necessary** for the following indications:

- A. A second ventilator to serve a different purpose from the first ventilator, based on medical needs. For example, two different types of ventilators are needed for each day, e.g., negative pressure ventilator with chest shell for one indication and a positive pressure ventilator with nasal mask the rest of the day;
- B. A back-up ventilator for one of the following:

1. Member/enrollee is confined to a wheelchair and requires a wheel-chair mounted ventilator during the day and another ventilator of the same type for use while in bed (unable to position the wheelchair-mounted ventilator close enough to the bed for use while sleeping). Without both pieces of equipment, member/enrollee may be prone to medical complications, unable to achieve appropriate medical outcomes, or may not be able to use the equipment effectively;
 2. Residence in remote areas with poor emergency access.
- V.** It is the policy of Louisiana Healthcare Connections that non-invasive home ventilators for overlap syndromes (presence of more than one condition, such as COPD and sleep apnea) require secondary review by a medical director.
- VI.** If the member needs the device long term, it appears that a purchase is more cost effective than a long term rental. This should be converted to a purchase if the provider agrees or sent to secondary review if they request long term rental.

Background

The term respiratory failure refers to the inability to adequately perform the fundamental functions of respiration, delivery of oxygen to the blood stream and removal of carbon dioxide. Respiratory failure has many causes and can be acute or chronic in nature. Typically, respiratory failure initially affects the ability to effectively move oxygen into the body, also known as oxygenation failure, or to eliminate carbon dioxide, also known as ventilatory failure.^{2,11}

Routine use of noninvasive ventilation has increased over the previous two decades, and as a result, noninvasive ventilation has become an essential tool in the management of acute and chronic respiratory failure in both the home and critical care settings.¹ Noninvasive ventilation offers increased flexibility and has become a valuable treatment option for patients with acidosis in moderate to severe respiratory distress and tachypnea with increased labored breathing due to COPD (chronic obstructive pulmonary disease) exacerbation.^{1,11}

Ventilatory support is achieved through a variety of interfaces such as oronasal mask, nasal mask, nasal prongs or full-face mask and by using a variety of ventilatory modes (e.g., volume ventilation, pressure support, cuirass ventilation, bi-level positive airway pressure [BiPAP], proportional-assist ventilation [PAV], continuous positive airway pressure [CPAP]). Oxygen is delivered via tubing through a positive pressure ventilator circuit and should be heated and humidified to improve tolerance and prevent mucosal dryness, a common side effect of prolonged noninvasive ventilation. The primary goals of home noninvasive ventilation are reduction of symptoms, improvement of quality of life, reduced readmission risk and reduction of mortality.^{1,2,3}

Invasive mechanical ventilation is primarily used to facilitate the exchange of oxygen and carbon dioxide, fully or partially, in patients with respiratory failure who no longer have the capacity to breathe spontaneously or whose ventilatory needs exceed their own ability to do so adequately. It is beneficial for protecting the airway of patients with a decreased level of consciousness, upper gastrointestinal hemorrhage, emesis, or other conditions with an increased risk of aspiration in whom noninvasive ventilation is contraindicated.^{12,13}

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2022, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only and may not support medical necessity. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
E0465	Home ventilator, any type, used with invasive interface, (e.g., tracheostomy tube)
E0466	Home ventilator, any type, used with noninvasive interface, (e.g., mask, chest shell)
E0467	Home ventilator, multi-function respiratory device, also performs any or all of the additional functions of oxygen concentration, drug nebulization, aspiration, and cough stimulation, includes all accessories, components and supplies for all functions

Reviews, Revisions, and Approvals	Revision Date	Approval Date
Converted corporate to local policy.	08/15/20	
References reviewed and updated. ICD-10 codes removed.	1/2022	
Changed name of policy to Invasive and Non-Invasive Home Ventilators. Added criteria i and ii to I.A.b. Removed from I.C.d: worsened greater than or equal to 7 mm Hg compared to the original result (see C.2). In section II, added 12-month certification after initial 3 month. Added HCPCS Code E0467. Removed invasive ventilator criteria from CP.MP.107 DME and placed in criteria. Description and background updated to include information re: invasive ventilators. References reviewed and updated.	10/2022	1/14/23
Annual review completed. Minor rewording with no clinical significance. Background updated with no clinical significance. References reviewed and updated.	07/23	9/25/23

References

1. SooHoo, G.W. Noninvasive ventilation. Medscape. <https://emedicine.medscape.com/article/304235-overview>. Updated June 18, 2020. Accessed April 20, 2023.

2. Hyzy RC, McSparron JI. Noninvasive ventilation in adults with acute respiratory failure: Practical aspects of initiation. UpToDate. www.uptodate.com. Updated March 28, 2023. Accessed April 20, 2023.
3. Breathing in America: Diseases, progress, and hope. American Thoracic Society. Published 2010.
4. Huttmann SE, Storre JH, Windisch W. Home mechanical ventilation: Invasive and noninvasive ventilation therapy for chronic respiratory failure. *Anaesthetist*. 2015;64(6):479 to 487. doi:10.1007/s00101-015-0049-z
5. Local Coverage Determination: Respiratory Assist Devices (L33800). Centers for Medicare and Medicaid Services Web site. <https://www.cms.gov/medicare-coverage-database/view/lcd.aspx?lcdid=33800&ver=26&bc=0>. Published October 1, 2015 (revised August 8, 2021). Accessed April 20, 2023.
6. Simonds AK. Home Mechanical Ventilation: An Overview. *Ann Am Thorac Soc*. 2016;13(11):2035 to 2044. doi:10.1513/AnnalsATS.201606-454FR
7. Gay PC. Nocturnal ventilator support in COPD. UpToDate. www.uptodate.com. Updated January 4, 2023. Accessed April 20, 2023.
8. National Coverage Determinations Manual (Internet-Only Manual, Publ. 100-03, Chapter 1, Part 4, Section 280.1). Centers for Medicare and Medicaid Services Web site. https://www.cms.gov/regulations-and-guidance/guidance/manuals/downloads/ncd103c1_part4.pdf. Published October 1, 2003. (revised March 9, 2023). Accessed April 20, 2023.
9. Hyzy RC, McSparron JI. Noninvasive ventilation in adults with acute respiratory failure: Benefits and contraindications. UpToDate. www.uptodate.com. August 31, 2021. Accessed April 20, 2023.
10. Martin TJ. Noninvasive positive airway pressure therapy for the obesity hypoventilation syndrome. UpToDate. www.uptodate.com. Updated May 20, 2022. Accessed April 20, 2023.
11. Feller-Kopman DJ, Schwartzstein RM. The evaluation, diagnosis, and treatment of the adult patient with acute hypercapnic respiratory failure. UpToDate. www.uptodate.com. Updated June 14, 2022. Accessed April 20, 2023.
12. Harmon EM. Acute respiratory distress syndrome (ARDS) treatment & management. Medscape. <https://emedicine.medscape.com/article/165139-treatment#d9>. Updated March 27, 2020. Accessed April 20, 2023.
13. Hyzy RC, McSparron JI. Overview of initiating invasive mechanical ventilation in adults in the intensive care unit. UpToDate. www.uptodate.com. Updated June 14, 2022. Accessed April 20, 2023.
14. Howell JD. Acute severe asthma exacerbations in children younger than 12 years: endotracheal intubation and mechanical ventilation. UpToDate. www.uptodate.com. Updated July 18, 2018. Accessed April 20, 2023.
15. Onweni C, Rashid S, Goswami R, et al. Cuirass ventilation: an alternative home-based modality for chronic respiratory failure. *Home Health Care Management & Practice*. 2020;32(1):40 to 44. doi:10.1177/1084822319875111
16. Waseem M. Pediatric pneumonia clinical presentation. Medscape. <https://emedicine.medscape.com/article/967822-clinical#b3>. Updated March 10, 2023. Accessed April 20, 2023.
17. Nagler JN. Noninvasive ventilation for acute and impending respiratory failure in children. UpToDate. www.uptodate.com. Updated June 29, 2022. Accessed April 20, 2023.

18. Hill NS, Kramer NR. Noninvasive ventilation in adults with chronic respiratory failure from neuromuscular and chest wall diseases: practical aspects of initiation. UpToDate. www.uptodate.com. Updated May 9, 2022. Accessed April 21, 2023.
19. Bach JR. Noninvasive ventilatory support and mechanical insufflation-exsufflation for patients with respiratory muscle dysfunction. UpToDate. www.uptodate.com. Updated March 2, 2023. Accessed April 21, 2023.
20. Hill NS, Kramer NR. Noninvasive ventilation in adults with chronic respiratory failure from neuromuscular and chest wall diseases: patient selection and alternative modes of ventilatory support. UpToDate. www.uptodate.com. Updated May 11, 2022. Accessed April 21, 2023.

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care and are solely responsible for the medical advice and treatment of members/enrollees. This clinical policy is not intended to recommend treatment for members/enrollees. Members/enrollees should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members/enrollees, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members/enrollees and their representatives agree to be bound by such terms and conditions by providing services to members/enrollees and/or submitting claims for payment for such services.

©2023 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademarks exclusively owned by Louisiana Healthcare Connections.