

## Clinical Policy: Antithymocyte Globulin (Atgam, Thymoglobulin)

Reference Number: LA.PHAR.506

Effective Date:

Last Review Date: 07.24.23

Line of Business: Medicaid

[Coding Implications](#)  
[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

**\*\*Please note: This policy is for medical benefit\*\***

### Description

Antithymocyte globulin (Thymoglobulin<sup>®</sup>, Atgam<sup>®</sup>) is an immunoglobulin G.

### FDA Approved Indication(s)

Atgam is indicated for:

- The management of allograft rejection in renal transplant patients; when administered with conventional therapy at the time of rejection, Atgam increases the frequency of resolution of the acute rejection episode.
- The treatment of moderate-to-severe aplastic anemia in patients unsuitable for bone marrow transplantation.

Limitation(s) of use: The usefulness of Atgam has not been demonstrated in patients with aplastic anemia who are suitable candidates for bone marrow transplantation or in patients with aplastic anemia secondary to neoplastic disease, storage disease, myelofibrosis, Fanconi's syndrome, or in patients known to have been exposed to myelotoxic agents or radiation.

Thymoglobulin is indicated for the prophylaxis and treatment of acute rejection in patients receiving a kidney transplant. Thymoglobulin is used in conjunction with concomitant immunosuppression.

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of Louisiana Healthcare Connections<sup>®</sup> that Atgam and Thymoglobulin are **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Kidney Transplant Rejection (must meet all):

1. Member has received or is scheduled for a kidney transplant;
2. If request is for prophylaxis of acute rejection, request is for Thymoglobulin;
3. Prescribed by or in consultation with a nephrologist, transplant specialist, or hematologist/oncologist;
4. Age  $\geq$  18 years;
5. Dose does not exceed one of the following (a or b):

## CLINICAL POLICY

### Antithymocyte Globulin

- a. For Atgam: 15 mg/kg per day;
- b. For Thymoglobulin: 1.5 mg/kg per day.

**Approval duration:**

7 days for Thymoglobulin for prophylaxis of acute rejection (7 doses)  
14 days for Thymoglobulin for treatment of acute rejection (14 doses)  
Up to 42 days for Atgam (21 doses)

**B. Aplastic Anemia (must meet all):**

1. Diagnosis of moderate to severe aplastic anemia;
2. Request is for Atgam;
3. Prescribed by or in consultation with a hematologist;
4. Age  $\geq$  18 years;
5. Prescribed in combination with cyclosporine;
6. Dose does not exceed 20 mg/kg per day.

**Approval duration:** Up to 42 days (21 doses)

**C. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

## II. Continued Therapy

**A. All Indications in Section I (must meet all):**

1. Currently receiving medication via Louisiana Healthcare Connections benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed (a or b):
  - a. For Atgam (i or ii):
    - i. For treatment of acute rejection: 15 mg/kg per day;
    - ii. For aplastic anemia: 20 mg/kg per day;
  - b. For Thymoglobulin for treatment or prophylaxis of acute rejection: 1.5 mg/kg per day.

**Approval duration:** Up to a total treatment duration of:

7 days for Thymoglobulin for prophylaxis of acute rejection (7 doses)  
14 days for Thymoglobulin for treatment of acute rejection (14 doses)  
42 days for Atgam (21 doses)

**B. Other diagnoses/indications (must meet 1 or 2):**

## CLINICAL POLICY

### Antithymocyte Globulin

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

#### III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – LA.PMN.53 for Medicaid, or evidence of coverage documents.

#### IV. Appendices/General Information

##### *Appendix A: Abbreviation/Acronym Key*

FDA: Food and Drug Administration

##### *Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
cyclosporine	<b>Aplastic Anemia</b> Adults: 12 mg/kg PO QD Children: 15 mg/kg PO QD	See dosing regimen

*Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.*

##### *Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s):
  - Atgam: patients with a history of a systemic reaction (e.g., anaphylactic reaction) during prior administration of Atgam or any other equine gamma globulin preparation
  - Thymoglobulin:
    - Patients with history of allergy or anaphylactic reaction to rabbit proteins or to any product excipients
    - Patients who have active acute or chronic infections that contraindicate any additional immunosuppression
- Boxed warning(s):
  - Atgam: anaphylaxis
  - Thymoglobulin: immunosuppression

##### *Appendix D: General Information*

## CLINICAL POLICY

### Antithymocyte Globulin

- The current standard first-line treatment for aplastic anemia is equine antithymocyte globulin (Atgam) combined with cyclosporine (off-label use).

#### V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Antithymocyte globulin (Atgam)	Aplastic anemia	10 to 20 mg/kg IV QD for 8 to 14 days. Additional alternate-day therapy up to a total of 21 doses may be given.	20 mg/kg/dose
Antithymocyte globulin (Atgam)	Treatment of acute renal transplant rejection	10 to 15 mg/kg IV QD for 14 days. Additional alternate-day therapy up to a total of 21 doses may be given.	15 mg/kg/dose
Antithymocyte globulin (Thymoglobulin)	Prophylaxis of acute renal transplant rejection	1.5 mg/kg IV QD for 4 to 7 days	1.5 mg/kg/dose
Antithymocyte globulin (Thymoglobulin)	Treatment of acute renal transplant rejection	1.5 mg/kg IV QD for 7 to 14 days	1.5 mg/kg/dose

#### VI. Product Availability

Drug Name	Availability
Antithymocyte globulin (Thymoglobulin)	Vial, powder for solution: 25 mg
Antithymocyte globulin (Atgam)	Ampule: 250 mg/5 mL

#### VII. References

- Thymoglobulin Prescribing Information. Cambridge, MA: Genzyme Corporation; March 2023. Available at: <http://products.sanofi.us/Thymoglobulin/Thymoglobulin.pdf>. Accessed July 24, 2023.
- Atgam Prescribing Information. New York, NY: Pfizer; May 2023. Available at: <http://labeling.pfizer.com/ShowLabeling.aspx?id=525>. Accessed July 24, 2023.
- Kidney Disease Improving Global Outcomes. KDIGO clinical practice guideline for the care of kidney transplant recipients. American Journal of Transplantation 2009; 9 (Suppl 3): S1-S155. doi: 10.1111/j.1600-6143.2009.02834.x
- Bia M, Adey DB, Bloon RD, Chan L, Kulkarni S, and Tomlanovich S. KDOQI US Commentary on the 2009 KDIGO clinical practice guideline for the care of kidney transplant recipients. Am J Kidneys Dis 2010;56:189-218.
- Schinstock CA, Mannon RB, Budde K, et al. Recommended treatment for antibody-mediated rejection after kidney transplantation: the 2019 expert consensus from the Transplantation Society Working Group. Transplantation May 2020;104(5):911-22.

## CLINICAL POLICY

### Antithymocyte Globulin

6. Cooper JE. Evaluation and treatment of acute kidney rejection in kidney allografts. CJASN March 2020;15:430-8.
7. Killick SB, Bown N, Cavenagh J, et al. Guidelines for the diagnosis and management of adult aplastic anaemia. Br J Haematol. 2016; 172:187-207.
8. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2019. Available at: <http://www.clinicalpharmacology-ip.com/>.

#### Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J7504	Lymphocyte immune globulin, antithymocyte globulin, equine, parenteral, 250 mg
J7511	Lymphocyte immune globulin, antithymocyte globulin, rabbit, parenteral, 25 mg

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Policy created	05.01.23	
References reviewed and updated.	07.24.23	

#### Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

## CLINICAL POLICY

### Antithymocyte Globulin

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2023 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.