

## Clinical Policy: Elotuzumab (Empliciti)

Reference Number: CP.PHAR.308

Effective Date: 02.01.17

Last Review Date: 11.23

Line of Business: Commercial, HIM, Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### Description

Elotuzumab (Empliciti<sup>®</sup>) is a SLAMF7-directed immunostimulatory antibody.

### FDA Approved Indication(s)

Empliciti is indicated in combination with:

- Lenalidomide and dexamethasone for the treatment of patients with multiple myeloma (MM) who have received one to three prior therapies
- Pomalidomide and dexamethasone for the treatment of adult patients with MM who have received at least two prior therapies including lenalidomide and a proteasome inhibitor

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Empliciti is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Multiple Myeloma (must meet all):

1. Diagnosis of MM;
  2. Prescribed by or in consultation with an oncologist;
  3. Age  $\geq$  18 years;
  4. Member has received  $\geq$  1 prior therapy (*see Appendix B for examples*);
  5. Empliciti is prescribed in combination with dexamethasone, and either Pomalyst<sup>®</sup>, lenalidomide, or bortezomib;\*
- \*Prior authorization may be required for Pomalyst, lenalidomide, and bortezomib.*
6. Request meets one of the following (a or b):\*
    - a. Dose does not exceed (i or ii):
      - i. With lenalidomide, both of the following (1 and 2):
        - 1) 10 mg/kg per week for the first two cycles (4 doses per 28-day cycle);
        - 2) 10 mg/kg per 2 weeks (2 doses per 28-day cycle) for subsequent cycles;
      - ii. With Pomalyst, both of the following (1 and 2):
        - 1) 10 mg/kg every week for the first 2 cycles (4 doses per 28-day cycle);
        - 2) 20 mg/kg every 4 weeks (1 dose per 28-day cycle) for subsequent cycles;
    - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN.*

**Approval duration: 6 months**

**B. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**II. Continued Therapy**

**A. Multiple Myeloma (must meet all):**

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Empliciti for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following\* (a or b):
  - a. New dose does not exceed (i or ii):
    - i. With lenalidomide, both of the following (1 and 2):
      - 1) 10 mg/kg per week for the first two cycles (4 doses per 28-day cycle);
      - 2) 10 mg/kg per 2 weeks (2 doses per 28-day cycle) for subsequent cycles;
    - ii. With Pomalyst, both of the following (1 and 2):
      - 1) 10 mg/kg every week for the first 2 cycles (4 doses per 28-day cycle);
      - 2) 20 mg/kg every 4 weeks (1 dose per 28-day cycle) for subsequent cycles;
  - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN.*

**Approval duration: 12 months**

**B. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or

- b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace and CP.PMN.53 for Medicaid or evidence of coverage documents.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

FDA: Food and Drug Administration

MM: multiple myeloma

NCCN: National Comprehensive Cancer Network

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

<b>Drug Name</b>	<b>Dosing Regimen</b>	<b>Dose Limit/ Maximum Dose</b>
bortezomib (Velcade)	<u>Empliciti in combination with Velcade and dexamethasone:</u> <ul style="list-style-type: none"> <li>• Regimens vary.</li> <li>• Per NCCN, the SC rather than IV bortezomib formulation is preferred. <i>An SC generic formulation is not available.</i></li> </ul>	Varies
lenalidomide (Revlimid)	<u>Empliciti in combination with Revlimid and dexamethasone:</u> Regimens vary.	
Pomalyst (pomalidomide)	<u>Empliciti in combination with Pomalyst and dexamethasone:</u> Regimens vary.	
Kyprolis <sup>®</sup> (carfilzomib), bortezomib (Velcade),	<u>Examples of primary therapy</u> <ul style="list-style-type: none"> <li>• Bortezomib/dexamethasone</li> <li>• Bortezomib/lenalidomide/dexamethasone</li> <li>• Bortezomib/cyclophosphamide/dexamethasone</li> </ul>	Varies

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
lenalidomide (Revlimid), cyclophosphamide, dexamethasone	<ul style="list-style-type: none"> <li>• Bortezomib/doxorubicin/dexamethasone</li> <li>• Bortezomib/thalidomide/dexamethasone</li> <li>• Carfilzomib/cyclophosphamide/dexamethasone</li> <li>• Carfilzomib/lenalidomide/dexamethasone</li> <li>• Cyclophosphamide/lenalidomide/dexamethasone</li> <li>• Daratumumab/lenalidomide/dexamethasone</li> <li>• Daratumumab/lenalidomide/bortezomib/dexamethasone</li> <li>• Daratumumab/carfilzomib/lenalidomide/dexamethasone</li> <li>• Daratumumab/cyclophosphamide/bortezomib/dexamethasone</li> <li>• Daratumumab/bortezomib/thalidomide/dexamethasone</li> <li>• Daratumumab/bortezomib/melphalan/prednisone</li> <li>• Dexamethasone/thalidomide/cisplatin/doxorubicin/cyclophosphamide/etoposide/bortezomib (VTD-PACE)</li> <li>• Ixazomib/cyclophosphamide/dexamethasone</li> <li>• Ixazomib/lenalidomide/dexamethasone</li> <li>• Lenalidomide/low-dose dexamethasone</li> </ul>	
Kyprolis (carfilzomib), bortezomib (Velcade), lenalidomide (Revlimid), Darzalex <sup>®</sup> (daratumumab), Ninlaro <sup>®</sup> (ixazomib), Pomalyst (pomalidomide), Empliciti <sup>®</sup> (elotuzumab), Thalomid <sup>®</sup> (thalidomide), bendamustine, cyclophosphamide, dexamethasone, Sarclisa <sup>®</sup>	<p><u>Examples of therapy for previously treated for relapsed or refractory disease:</u></p> <ul style="list-style-type: none"> <li>• Bendamustine</li> <li>• Bendamustine/bortezomib/dexamethasone</li> <li>• Bendamustine/lenalidomide/dexamethasone</li> <li>• Bendamustine/carfilzomib/dexamethasone</li> <li>• Bortezomib/dexamethasone</li> <li>• Bortezomib/lenalidomide/dexamethasone</li> <li>• Bortezomib/liposomal doxorubicin/dexamethasone</li> <li>• Bortezomib/cyclophosphamide/dexamethasone</li> <li>• Carfilzomib/cyclophosphamide/dexamethasone</li> <li>• Carfilzomib/dexamethasone</li> <li>• Carfilzomib/lenalidomide/dexamethasone</li> <li>• Carfilzomib/cyclophosphamide/dexamethasone</li> <li>• Carfilzomib/cyclophosphamide/thalidomide/dexamethasone</li> <li>• Cyclophosphamide/lenalidomide/dexamethasone</li> <li>• Cyclophosphamide</li> <li>• Daratumumab</li> </ul>	Varies

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
(istatuximab-irfc), Xpovio <sup>®</sup> (selinexor)	<ul style="list-style-type: none"> <li>• Daratumumab/bortezomib/dexamethasone</li> <li>• Daratumumab/carfilzomib/dexamethasone</li> <li>• Daratumumab/cyclophosphamide/bortezomib/dexamethasone</li> <li>• Daratumumab/lenalidomide/dexamethasone</li> <li>• Daratumumab/pomalidomide/dexamethasone</li> <li>• Dexamethasone/cyclophosphamide/etoposide/cisplatin</li> <li>• Dexamethasone/thalidomide/cisplatin/doxorubicin/cyclophosphamide/etoposide/ +/- bortezomib</li> <li>• Elotuzumab/lenalidomide/dexamethasone</li> <li>• Elotuzumab/bortezomib/dexamethasone</li> <li>• Elotuzumab/pomalidomide/dexamethasone</li> <li>• Istatuximab-irfc/carfilzomib/dexamethasone</li> <li>• Ixazomib/cyclophosphamide/dexamethasone</li> <li>• Ixazomib/lenalidomide/dexamethasone</li> <li>• Ixazomib/pomalidomide/dexamethasone</li> <li>• Isatuximab-irfc/pomalidomide/dexamethasone</li> <li>• Lenalidomide/dexamethasone</li> <li>• Pomalidomide/bortezomib/dexamethasone</li> <li>• Pomalidomide/carfilzomib/dexamethasone</li> <li>• Pomalidomide/cyclophosphamide/dexamethasone</li> <li>• Pomalidomide/dexamethasone</li> <li>• Selinexor/bortezomib/dexamethasone</li> <li>• Selinexor/carfilzomib/dexamethasone</li> <li>• Selinexor/daratumumab/dexamethasone</li> <li>• Selinexor/pomalidomide/dexamethasone</li> <li>• Venetoclax/dexamethasone</li> <li>• Ideocabtagene vicleucel</li> <li>• Ciltacabtagene autoleucel</li> <li>• Teclistamab-cqyv</li> <li>• Benlantamab mafodotin-blmf</li> </ul>	

*Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.*

*Appendix C: Contraindications/Black Box Warnings*

None reported

**V. Dosage and Administration**

Indication	Dosing Regimen	Maximum Dose
MM	<p><u>Cycles one and two:</u></p> <ul style="list-style-type: none"> <li>• Empliciti: 10 mg/kg IV once weekly on cycles 1 and 2 (on days 1, 8, 15, and 22),</li> <li>• Dexamethasone: 28 mg PO between 3 and 24 hours before Empliciti plus 8 mg IV between 45 and 90 minutes before Empliciti</li> <li>• Lenalidomide: 25 mg PO QD x 21 days of a 28-day cycle</li> </ul> <p>OR</p> <ul style="list-style-type: none"> <li>• Pomalidomide: 4 mg PO QD x 21 days of a 28-day cycle</li> </ul> <p><u>Cycles three and beyond:</u></p> <ul style="list-style-type: none"> <li>• Empliciti:               <ul style="list-style-type: none"> <li>○ With lenalidomide: 10 mg/kg IV once every 2 weeks (on days 1 and 15)</li> <li>○ With pomalidomide: 20 mg/kg IV once every 4 weeks</li> </ul> </li> <li>• Dexamethasone: Administer as for cycles one and two and on the days Empliciti is not given (days 8 and 22), give 40 mg PO QD if 75 years or younger OR 20 mg PO QD if older than 75 years</li> <li>• Lenalidomide: 25 mg PO QD x 21 days of a 28-day cycle</li> </ul> <p>OR</p> <ul style="list-style-type: none"> <li>• Pomalidomide: 4 mg PO QD x 21 days of a 28-day cycle</li> </ul>	<p>With lenalidomide: 10 mg/kg</p> <p>With pomalidomide: 20 mg/kg</p>

**VI. Product Availability**

Single-dose vial: 300 mg, 400 mg

**VII. References**

1. Empliciti Prescribing Information. Princeton, NJ: Bristol-Myers Squibb; March 2022. Available at: [https://packageinserts.bms.com/pi/pi\\_empliciti.pdf](https://packageinserts.bms.com/pi/pi_empliciti.pdf). Accessed August 5, 2023.
2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: [http://www.nccn.org/professionals/drug\\_compendium](http://www.nccn.org/professionals/drug_compendium). Accessed August 5, 2023.
3. National Comprehensive Cancer Network. Multiple Myeloma Version 3.2023. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/myeloma.pdf](https://www.nccn.org/professionals/physician_gls/pdf/myeloma.pdf). Accessed August 5, 2023.

**Coding Implications**

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J9176	Injection, elotuzumab, 1 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
RT4: added newly FDA-approved use with pomalidomide for MM; references reviewed and updated.	11.27.18	02.19
4Q 2019 annual review: FDA/NCCN dosing requirement added; references reviewed and updated.	08.20.19	11.19
4Q 2020 annual review: added Commercial line of business, modified HIM-Medical Benefit to HIM line of business; references reviewed and updated.	08.11.20	11.20
4Q 2021 annual review: updated Appendix B Therapeutic Alternatives; modified reference from HIM.PHAR.21 to HIM.PA.154; references reviewed and updated.	08.11.21	11.21
4Q 2022 annual review: no significant changes; updated Appendix B per NCCN MM guidelines for primary therapy and therapy for previously treated MM; references reviewed and updated. Template changes applied to other diagnoses/indications.	07.28.22	11.22
4Q 2023 annual review: no significant changes; updated Appendix B with examples of previously treated regimens per current NCCN Multiple Myeloma guidelines; references reviewed and updated.	08.05.23	11.23

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note:**

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2017 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.