

# Clinical Policy: Attention Deficit Hyperactivity Disorder Assessment and Treatment

Reference Number: LA.CP.BH.124

Date of Last Revision: 3/25

Coding Implications
Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

#### **Description**

Attention deficit hyperactivity disorder (ADHD) is one of the most common neurobehavioral disorders in children, with an increasing prevalence of diagnosis in adults. ADHD affects the cognitive, academic, emotional, and social well-being of individuals and can persist throughout life. While there is no single test to diagnose ADHD, a clinical assessment based on defined clinical parameters establishes criteria for diagnosis in children and adults.

#### Policy/Criteria

I. It is the policy of Louisiana Healthcare Connections that the following services are **medically necessary** when requested for the assessment and treatment of attention deficit hyperactivity disorder (ADHD):

#### A. Assessment

- 1. Complete medical evaluation with history and physical examination;
- 2. Parent/child interview or patient interview, if adult, to obtain information listed in Diagnostic and Statistical Manual of Mental Health Disorders, Fifth Edition (DSM-5 TR):
- 3. Collection of collateral information, if available, such as the Vanderbilt or Conners assessment;
- 4. Complete psychiatric evaluation or other services provided by a psychiatrist, psychologist, or other behavioral health professional;
- 5. Laboratory evaluation prior to stimulant medication therapy, including any of the following:
  - a. Complete blood count;
  - b. Liver function tests;
  - c. Toxicology screen if drug use is suspected;
  - d. Cardiac evaluation and screening. Electrocardiogram (ECG), if clinically indicated (e.g., family or personal history of cardiovascular disease or those with congenital heart disease);
- 6. Measurement of thyroid hormone levels if patient exhibits clinical manifestations of hyperthyroidism;
- 7. Assessment of comorbid behavioral health and/or medical diagnoses and associated symptoms;
- 8. When not otherwise excluded, other services for the assessment of ADHD to meet the DSM-5 TR criteria.

#### B. Treatment:

- 1. Pharmacotherapy;
- 2. Behavioral modification;



- 3. Treatment of comorbid behavioral health and/or medical diagnoses and associated symptoms;
- 4. When not otherwise excluded, other services for the treatment of ADHD;
- 5. Ongoing assessment and application of standardized scales to assess treatment benefit.
- 6. Treatment of family, guardian(s) and/or custodian. Also, parenting therapy has been shown to be an effective treatment in addressing family issues in children with ADHD. Evaluations should be conducted on family members who appear to be contributing to family disorder or on family members negatively impacted by the member's ADHD.
- II. It is the policy of Louisiana Healthcare Connections that there is insufficient evidence to support the following for the assessment or treatment of ADHD (may not be all-inclusive):
  - **A.** Assessment:
    - 1. Actigraphy;
    - 2. Acoustic reflex testing;
    - 3. AFF2 gene testing;
    - 4. Assessment of serum lipid profiles;
    - 5. Computerized electroencephalogram (EEG) (e.g., brain mapping, neurometrics, or quantitative electroencephalography [qEEG], Neuropsychiatric EEG-Based Assessment Aid [NEBA] System);
    - 6. Computerized tests of attention and vigilance;
    - 7. Education and achievement testing;
    - 8. Electronystagmography in the absence of symptoms of vertigo or balance dysfunction;
    - 9. Evaluation of iron status (e.g. measurement of serum iron and ferritin levels);
    - 10. Event-related potentials;
    - 11. Functional near-infrared spectroscopy;
    - 12. Hair analysis:
    - 13. IgG blood tests;
    - 14. Measurement of peripheral brain-derived neurotrophic factor;
    - 15. Measurement of zinc;
    - 16. Neuroimaging (e.g., CT [computed tomography], CAT [computerized axial tomography], MRI [magnetic resonance imaging], including diffusion tensor imaging), MRS (magnetic resonance spectroscopy), PET (positron emission tomography), and SPECT (single-photon emission computerized tomography), functional brain mapping;
    - 17. Neuropsychiatric EEG-based assessment aid system;
    - 18. Pharmacogenetic tools (vascular flow brain imaging);
    - 19. Otoacoustic emissions in the absence of signs of hearing loss;
    - 20. Quotient ADHD system/test;
    - 21. Synaptosomal-associated protein (SNAP) 25 gene polymorphisms testing;
    - 22. Transcranial magnetic stimulation—evoked measures (e.g., short-interval cortical inhibition in motor cortex) as a marker of ADHD symptoms;
    - 23. Measures of thyroid hormones unless the individual exhibits clinical manifestations of hyperthyroidism (e.g. (modest acceleration of linear growth andepiphyseal maturation, weight loss or failure to gain weight, excessive retraction of the eyelids



causing lid lag and stare, diffuse goiter, tachycardia and increased cardiac output, increased gastrointestinal motility, tremor, hyperreflexia);

24. Tympanometry in the absence of hearing loss.

#### **B.** Treatment:

- 1. Acupuncture/acupressure;
- 2. Application of: hot or cold packs, traction, mechanical, electrical stimulation (unattended), vasopneumatic devices, paraffin bath, whirlpool, diathermy (eg, microwave), infrared, ultraviolet, electrical stimulation (manual), iontophoresis, contrast baths, ultrasound, hubbard tank;
- 3. Anti-candida albicans medication;
- 4. Anti-fungal medication;
- 5. Anti-motion sickness medication;
- 6. Auditory Integration Therapy;
- 7. Applied kinesiology;
- 8. Brain integration;
- 9. Cannabinoids and cannabinoid products;
- 10. Chelation;
- 11. Chiropractic manipulation;
- 12. Cognitive behavior modification;
- 13. Cognitive rehabilitation;
- 14. Cognitive training;
- 15. Computerized training on working memory;
- 16. Deep pressure sensory vest;
- 17. Dietary counseling and treatments (i.e., Feingold diet);
- 18. Dore program/dyslexia–dyspraxia attention treatment (DDAT);
- 19. EEG biofeedback/neurofeedback;
- 20. External trigeminal nerve stimulation (eTNS);
- 21. Herbal remedies;
- 22. Homeopathy;
- 23. Intensive behavioral intervention programs;
- 24. Megavitamin therapy;
- 25. Metronome training;
- 26. Mindfulness;
- 27. Mineral supplementation;
- 28. Music therapy;
- 29. Optometric vision training;
- 30. Psychopharmaceuticals (lithium, benzodiazepines, and selective serotonin reuptake inhibitors, unless the patient also exhibits anxiety and depression);
- 31. Reboxetine;
- 32. Sensory integration therapy;
- 33. The Good Vibrations device;
- 34. The Neuro Emotional Technique;
- 35. Therapeutic eurythmy (movement therapy);



- 36. Transcranial magnetic stimulation/cranial electric stimulation;
- 37. Video game-based therapeutic interventions (e.g., EndeavorRx, AKL-T01)
- 38. Vision therapy;
- 39. Yoga.

III. It is the policy of Louisiana Healthcare Connections that interventions that are strictly educational in nature (e.g., classroom environmental manipulation, academic skills training) are not medically necessary as they are not considered medical interventions.

#### **Background**

ADHD (Attention Deficit Hyperactivity Disorder) is one of the most commonly diagnosed neurodevelopmental disorders in children and adolescents and is increasingly being diagnosed in adults.<sup>5</sup> The main characteristics of ADHD are symptoms of inattention, hyperactivity, and impulsivity that have continued for at least six months and are maladaptive and inconsistent with development level.<sup>1</sup> There is no single genetic or behavioral test to diagnose ADHD. Instead, a clinical diagnosis based on the *Diagnostic and Statistical Manual of Mental Disorders-5* (DSM-5) criteria is applicable for both children and adults.<sup>2</sup> Due to the prevalence of children and adolescents with an ADHD diagnosis, treatment of ADHD is often managed in the primary care setting, and evidence supports that appropriate diagnosis can be accomplished in this setting.<sup>5</sup> However, primary care providers should refer children to a specialist for complex ADHD symptoms.<sup>16</sup> Some of the more common comorbid disorders include anxiety, autism spectrum disorder, depression, disruptive behavior disorders, substance use disorders and Tic disorders.<sup>3,16</sup> Suggested first line treatment for adults with ADHD is medication rather than cognitive-behavioral therapy (CBT).<sup>18</sup>

#### American Academy of Pediatrics (AAP)

In 2011, the AAP published a clinical practice guideline to clarify the diagnosis, evaluation, and treatment parameters of ADHD with an update in 2019.<sup>4</sup> This guideline expanded the age range of children to include preschool aged children (4 to 6 years of age) and adolescents (12 to 18 years of age), and suggests an expanded scope for behavioral interventions.<sup>4</sup> The evaluation of comorbid conditions, including behavioral, emotional, developmental, and physical, that might coexist with ADHD must also be considered. 4,5 Most children and adolescents diagnosed with ADHD also meet diagnostic criteria for other behavioral health conditions. In some situations, the presence of a comorbid diagnosis will alter the course of ADHD treatment. Additionally, when an adolescent receives a new diagnosis of ADHD, an assessment for substance use, anxiety, depression, and learning disorders should also be conducted, as these are common comorbid conditions that may alter the treatment approach of the adolescent population.<sup>5</sup> Similar clinical recommendations have been made by various organizations for adults, including the Canadian ADHD Resource Alliance, the American Academy of the Child and Adolescent Psychiatry, the National Institutes of Health, and the British Association for Psyschopharmacology. 5 Pharmacotherapy can provide a way to manage ADHD symptoms and improve quality of life.

In 2020, The Society for Developmental and Behavioral Pediatrics (SDBP) published Clinical Practice Guideline for the Assessment and Treatment of Children and Adolescents with Complex



Attention-Deficit/Hyperactivity Disorder and Process of Care Algorithms (POCA) that are meant to be used as companion documents to the published guidelines. The algorithms include suggested steps in the treatment of complex ADHD and key concepts include focus on functional impairment to improve long-term outcomes, psychosocial treatment as foundational in the treatment of complex ADHD, shared decision making, interprofessional care, using mental health diagnostic assessment and testing appropriately, identifying and treating impairments caused by coexisting conditions, and a lifelong perspective. These algorithms are based on expert consensus, and review of existing publications and practice guidelines and are meant to improve the care that children and adolescents with complex ADHD receive.

Stimulants and non-stimulants are common examples of medications prescribed to treat ADHD. A systemic review of sixteen randomized clinical trials and one meta-analysis that involved 2668 participants and evaluated pharmacological and psychosocial treatments of ADHD in adolescents 12 to 18 years of age was completed. The findings demonstrated that extended-release methylphenidate and amphetamine formulations, atomoxetine, and extended-release guanfacine led to clinically significant symptom reduction. Nonstimulants are not approved by the FDA for use in preschool-aged children. There is strong evidence for stimulant medications and significant evidence, but less strong, for atomoxetine, extended release guanfacine, and extended-release clonidine. Due to the lack of significant studies in school-aged children for nonstimulant medication and dextroamphetamine, methylphenidate is recommended as the first line of pharmacologic treatment for this population.<sup>5</sup> Findings from clinical trials studying adults with noncomorbid ADHD suggest amphetamines as first-line treatment when compared to other medications or cognitive-behavioral therapy (CBT). 18 Methylphenidate is noted as the first option of treatment for adults with moderate or severe ADHD; however, the evidence on the effects of immediate-release (IR) methylphenidate is limited and controversial in the treatment of the adult population.<sup>17</sup>

The AAP (American Academy of Pediatrics) has established recommendations regarding treatment modalities based on age. It is recommended that preschool children (4 to 6 years of age) are first prescribed evidence-based behavioral Parent Training in Behavior Management (PTBM) and/or classroom interventions. If these methods are not effective, Methylphenidate can be considered. For elementary and middle school children (6 to 12 years of age), a combination of FDA approved medications for ADHD and PTBM and classroom interventions should be prescribed. Educational interventions and supports, including an Individualized Education Program (IEP) are a vital part of treatment. Adolescents (12 to 18 years of age) should be treated with FDA approved medications in conjunction with evidence-based training or behavioral interventions. Educational interventions and supports are also an important aspect of treatment in this age group and can include an IEP or 504 plan. Additionally, planning for adulthood is an important component of the chronic care model for ADHD.<sup>5</sup>

The AAP also recognizes psychosocial treatments as effective for the treatment of ADHD. These treatments may include behavioral therapy and training interventions. Behavioral therapy can help adults (parents and school staff) to learn how to respond effectively and prevent certain behaviors, such as interrupting, aggression, non-compliance with requests, and non-completion of tasks. Skill development is targeted in training interventions and include repeated practice and



performance feedback. The effectiveness of certain training interventions, such as social skills training, is not supported by research.<sup>5</sup>

While the pathogenesis of ADHD is unknown, the clinical impairments in neurobehavioral and neurodevelopmental functioning pathways elicit deficiencies in vigilance, perceptual-motor speed, working memory, verbal learning, and response inhibition.<sup>2</sup> Consequently, ADHD affects the cognitive, academic, emotional, and social wellbeing of individuals and can persist throughout life. ADHD is a chronic condition and children and adolescents with ADHD should be managed in the same way those with special health care needs would be managed. Principles of the chronic care model and the medical home should be followed.<sup>5</sup>

#### **Coding Implications**

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2023, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only and may not support medical necessity. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT codes considered not medically necessary when billed with a sole diagnosis of ADHD

<b>CPT</b> ®	Description			
Codes				
70450	Computed tomography, head or brain; without contrast material			
70460	Computed tomography, head or brain; with contrast material(s)			
70470	Computed tomography, head or brain; without contrast material, followed			
70496	by contrast material(s) and further sections  Computed tomographic angiography, head, with contrast material(s), including noncontrast images, if performed, and image postprocessing			
70544	Magnetic resonance angiography, head; without contrast material(s)			
70545	Magnetic resonance angiography, head; with contrast material(s)			
70546	Magnetic resonance angiography, head; without contrast material(s), followed by contrast material(s) and further sequences			
70551	Magnetic resonance (eg, proton) imaging, brain (including brain stem); without contrast material			
70552	Magnetic resonance (eg, proton) imaging, brain (including brain stem); with contrast material(s)			
70553	Magnetic resonance (eg, proton) imaging, brain (including brain stem); without contrast material, followed by contrast material(s) and further sequences			
70554	Magnetic resonance imaging, brain, functional MRI; including test selection and administration of repetitive body part movement and/or visual stimulation, not requiring physician or psychologist administration			



CPT®	Description			
Codes	•			
70555	Magnetic resonance imaging, brain, functional MRI; requiring physician or			
	psychologist administration of entire neurofunctional testing			
76390	Magnetic resonance spectroscopy			
78600	Brain imaging, less than 4 static views;			
78601	Brain imaging, less than 4 static views; with vascular flow			
78605	Brain imaging, minimum 4 static views;			
78606	Brain imaging, minimum 4 static views; with vascular flow			
78608	Brain imaging, positron emission tomography (PET); metabolic evaluation.			
78609	Brain imaging, positron emission tomography (PET); perfusion evaluation			
78610	Brain imaging, vascular flow only			
78803	Radiopharmaceutical localization of tumor, inflammatory process or			
	distribution of radiopharmaceutical agent(s) (includes vascular flow and			
	blood pool imaging, when performed); tomographic (SPECT), single area			
	(eg, head, neck, chest, pelvis), single day imaging			
80061	Lipid panel This panel must include the following: Cholesterol, serum, total			
	(82465) Lipoprotein, direct measurement, high density cholesterol (HDL			
	cholesterol) (83718) Triglycerides (84478)			
81171	AFF2 (AF4/FMR2 family, member 2 [FMR2]) (eg, fragile X mental			
	retardation 2 [FRAXE]) gene analysis; evaluation to detect abnormal (eg,			
	expanded) alleles			
81172	AFF2 (AF4/FMR2 family, member 2 [FMR2]) (eg, fragile X mental			
	retardation 2 [FRAXE]) gene analysis; characterization of alleles (eg,			
	expanded size and methylation status)			
81229	Cytogenomic (genome-wide) analysis for constitutional chromosomal			
	abnormalities; interrogation of genomic regions for copy number and single			
	nucleotide polymorphism (SNP) variants, comparative genomic			
	hybridization (CGH) microarray analysis			
82365	Calculus; Infrared spectroscopy			
82465	Cholesterol, serum or whole blood, total			
82728	Ferritin			
82784	Gammaglobulin (immunoglobulin); IgA, IgD, IgG, IgM, each			
82787	Gammaglobulin (immunoglobulin); immunoglobulin subclasses (eg, IgG1,			
	2, 3, or 4), each			
83540	Iron			
83550	Iron binding capacity			
83718	Lipoprotein, direct measurement; high density cholesterol (HDL cholesterol)			
83719	Lipoprotein, direct measurement; VLDL cholesterol			
83721	Lipoprotein, direct measurement; LDL cholesterol			
83722	Lipoprotein, direct measurement; small dense LDL cholesterol			
84436	Thyroxine; total			
84437	Thyroxine; requiring elution (eg, neonatal)			
84439	Thyroxine; free			
84442	Thyroxine binding globulin (TBG)			



<b>CPT</b> ®	Description			
Codes				
84443	Thyroid stimulating hormone (TSH)			
84445	Thyroid stimulating immune globulins (TSI)			
84478	Triglycerides			
84479	Thyroid hormone (T3 or T4) uptake or thyroid hormone binding ratio			
	(THBR)			
84481	Triiodothyronine T3; free			
84630	Zinc			
86001	Allergen specific IgG quantitative or semiquantitative, each allergen			
92065	Orthoptic training performed by a physician or other qualified health care			
	professional			
90867	Therapeutic repetitive transcranial magnetic stimulation (TMS) treatment;			
	initial, including cortical mapping, motor threshold determination, delivery			
	and management			
90868	Therapeutic repetitive transcranial magnetic stimulation (TMS) treatment;			
	subsequent delivery and management, per session			
90869 Therapeutic repetitive transcranial magnetic stimulation (TMS) to				
	subsequent motor threshold re-determination with delivery and management			
90901	Biofeedback training by any modality			
92540	Basic vestibular evaluation, includes spontaneous nystagmus test with			
	eccentric gaze fixation nystagmus, with recording, positional nystagmus test,			
	minimum of 4 positions, with recording, optokinetic nystagmus test,			
	bidirectional foveal and peripheral stimulation, with recording, and			
	oscillating tracking test, with recording			
92541	Spontaneous nystagmus test, including gaze and fixation nystagmus, with			
	recording			
92542	Positional nystagmus test, minimum of 4 positions, with recording			
92544	Optokinetic nystagmus test, bidirectional, foveal or peripheral stimulation,			
	with recording			
92547	Use of vertical electrodes (List separately in addition to code for primary			
00770	procedure)			
92550	Tympanometry and reflex threshold measurements			
92558	Evoked otoacoustic emissions, screening (qualitative measurement of			
	distortion product or transient evoked otoacoustic emissions), automated			
02565	analysis			
92567	Tympanometry (impedance testing)			
92568	Acoustic reflex testing, threshold			
92570	Acoustic immittance testing, includes tympanometry (impedance testing),			
02597	acoustic reflex threshold testing, and acoustic reflex decay testing			
92587	Distortion product evoked otoacoustic emissions; limited evaluation (to			
	confirm the presence or absence of hearing disorder, 3 to 6 frequencies) or			
02500	transient evoked otoacoustic emissions, with interpretation and report			
92588	Distortion product evoked otoacoustic emissions; comprehensive diagnostic			
	evaluation (quantitative analysis of outer hair cell function by cochlear			
	mapping, minimum of 12 frequencies), with interpretation and report			



<b>CPT</b> ®	Description			
Codes				
	Auditory evoked potentials; screening of auditory potential with broadband			
92650	stimuli, automated analysis			
92651	Auditory evoked potentials; for hearing status determination, broadband			
	stimuli, with interpretation and report			
92652	Auditory evoked potentials; for threshold estimation at multiple frequencies,			
	with interpretation and report			
92653	Auditory evoked potentials; neurodiagnostic, with interpretation and report			
93000	Electrocardiogram, routine ECG with at least 12 leads; with interpretation			
	and report			
93005	Electrocardiogram, routine ECG with at least 12 leads; tracing only, without			
	interpretation and report			
93010	Electrocardiogram, routine ECG with at least 12 leads; interpretation and			
	report only			
95803	Actigraphy testing recording, analysis, interpretation, and report (minimum			
	of 72 hours to 14 consecutive days of recording)			
95812	Electroencephalogram (EEG) extended monitoring; 41 to 60 minutes			
95813	Electroencephalogram (EEG) extended monitoring; 61 to 119 minutes			
95816	Electroencephalogram (EEG); including recording awake and drowsy			
95819	Electroencephalogram (EEG); including recording awake and asleep			
95705	Electroencephalogram (EEG), without video, review of data, technical			
	description by EEG technologist, 2 to 12 hours; unmonitored			
95706	Electroencephalogram (EEG), without video, review of data, technical			
	description by EEG technologist, 2 to 12 hours; with intermittent monitoring			
	and maintenance			
95707	Electroencephalogram (EEG), without video, review of data, technical			
	description by EEG technologist, 2 to 12 hours; with continuous, real-time			
	monitoring and maintenance			
95708 Electroencephalogram (EEG), without video, review of data, tech				
	description by EEG technologist, each increment of 12 to 26 hours;			
	unmonitored			
95709	Electroencephalogram (EEG), without video, review of data, technical			
	description by EEG technologist, each increment of 12 to 26 hours; with			
	intermittent monitoring and maintenance			
95710	Electroencephalogram (EEG), without video, review of data, technical			
	description by EEG technologist, each increment of 12 to 26 hours; with			
	continuous, real-time monitoring and maintenance			
95711	Electroencephalogram with video (VEEG), review of data, technical			
0.5.5.1.5	description by EEG technologist, 2 to 12 hours; unmonitored			
95712	Electroencephalogram with video (VEEG), review of data, technical			
	description by EEG technologist, 2 to 12 hours; with intermittent monitoring			
	and maintenance			
95713	Electroencephalogram with video (VEEG), review of data, technical			
	description by EEG technologist, 2 to 12 hours; with continuous, real-time			
	monitoring and maintenance			



<b>CPT</b> ®	Description			
Codes	Description .			
95714	Electroencephalogram with video (VEEG), review of data, technical description by EEG technologist, each increment of 12 to 26 hours; with continuous, real-time monitoring and maintenance			
95715	Electroencephalogram with video (VEEG), review of data, technical description by EEG technologist, each increment of 12 to 26 hours; with intermittent monitoring and maintenance			
95716	Electroencephalogram with video (VEEG), review of data, technical description by EEG technologist, each increment of 12 to 26 hours; with continuous, real-time monitoring and maintenance			
95717	Electroencephalogram (EEG), continuous recording, physician or other qualified health care professional review of recorded events, analysis of spike and seizure detection, interpretation and report, 2 to 12 hours of EEG recording; without video			
95718	Electroencephalogram (EEG), continuous recording, physician or other qualified health care professional review of recorded events, analysis of spike and seizure detection, interpretation and report, 2 to 12 hours of EEG recording; with video (VEEG)			
95719	Electroencephalogram (EEG), continuous recording, physician or other qualified health care professional review of recorded events, analysis of spike and seizure detection, each increment of greater than 12 hours, up to 26 hours of EEG recording, interpretation and report after each 24 hour period; without video			
95720	Electroencephalogram (EEG), continuous recording, physician or other qualified health care professional review of recorded events, analysis of spike and seizure detection, each increment of greater than 12 hours, up to 26 hours of EEG recording, interpretation and report after each 24 hour period; with video (VEEG)			
95721	Electroencephalogram (EEG), continuous recording, physician or other qualified health care professional review of recorded events, analysis of spike and seizure detection, interpretation, and summary report, complete study; greater than 36 hours, up to 60 hours of EEG recording, without video			
95722	Electroencephalogram (EEG), continuous recording, physician or other qualified health care professional review of recorded events, analysis of spike and seizure detection, interpretation, and summary report, complete study; greater than 36 hours, up to 60 hours of EEG recording, with video (VEEG)			
95723	Electroencephalogram (EEG), continuous recording, physician or other qualified health care professional review of recorded events, analysis of spike and seizure detection, interpretation, and summary report, complete study; greater than 60 hours, up to 84 hours of EEG recording, without video			
95724	Electroencephalogram (EEG), continuous recording, physician or other qualified health care professional review of recorded events, analysis of spike and seizure detection, interpretation, and summary report, complete			



CPT®	Description			
Codes				
	study; greater than 60 hours, up to 84 hours of EEG recording, with video (VEEG)			
95725	Electroencephalogram (EEG), continuous recording, physician or other qualified health care professional review of recorded events, analysis of spike and seizure detection, interpretation, and summary report, complete study; greater than 84 hours of EEG recording, without video			
95726	Electroencephalogram (EEG), continuous recording, physician or other qualified health care professional review of recorded events, analysis of spike and seizure detection, interpretation, and summary report, complete study; greater than 84 hours of EEG recording, with video (VEEG)			
95925	Short-latency somatosensory evoked potential study, stimulation of any/all peripheral nerves or skin sites, recording from the central nervous system; in upper limbs			
95926	Short latency somatosensory evoked potential study, stimulation of any/all peripheral nerves or skin sites, recording from the central nervous system; in lower limbs			
95927	Short latency somatosensory evoked potential study, stimulation of any/all peripheral nerves or skin sites, recording from the central nervous system; in the trunk or head			
95928	Central motor evoked potential study (transcranial motor stimulation); upper limbs			
95929	Central motor evoked potential study (transcranial motor stimulation); lower limbs			
95930	Visual evoked potential (VEP), checkerboard or flash testing, central nervous system except glaucoma, with interpretation and report			
95933	Orbicularis oculi (blink) reflex, by electrodiagnostic testing			
95937	Neuromuscular junction testing (repetitive stimulation paired stimuli), each nerve, any 1 method			
95938	Short latency somatosensory evoked potential study, stimulation of any/all peripheral nerves or skin sites, recording from the central nervous system; in upper and lower limbs			
95939	Central motor evoked potential study (transcranial motor stimulation); in upper and lower limbs			
95954	Pharmacological or physical activation requiring physician or other qualified health care professional attendance during EEG recording of activation phase (eg, thiopental activation test)			
95957	Digital analysis of electroencephalogram (EEG) (eg, for epileptic spike analysis)			
96020	Neurofunctional testing selection and administration during noninvasive imaging functional brain mapping, with test administered entirely by a physician or other qualified health care professional (ie, psychologist), with review of test results and report			
96116	Neurobehavioral status exam (clinical assessment of thinking, reasoning and judgment, eg, acquired knowledge, attention, language, memory, planning			



CPT®	Description			
Codes	Description			
Coucs	and problem solving, and visual spatial abilities), by physician or other			
	qualified health care professional, both face-to-face time with the patient and			
	time interpreting test results and preparing the report, first hour			
96121	Neurobehavioral status exam (clinical assessment of thinking, reasoning and			
	judgment, [eg, acquired knowledge, attention, language, memory, planning			
	and problem solving, and visual spatial abilities]), by physician or other			
	qualified health care professional, both face-to-face time with the patient and			
	time interpreting test results and preparing the report; each additional hour			
96365	Intravenous infusion, for therapy, prophylaxis, or diagnosis (specify			
	substance or drug); initial, up to 1 hour			
96366	Intravenous infusion, for therapy, prophylaxis, or diagnosis (specify			
	substance or drug); each additional hour (List separately in addition to code			
	for primary procedure)			
96367	Intravenous infusion, for therapy, prophylaxis, or diagnosis (specify			
	substance or drug); additional sequential infusion of a new drug/substance,			
	up to 1 hour (List separately in addition to code for primary procedure)			
96902 Microscopic examination of hairs plucked or clipped by the examina				
	(excluding hair collected by the patient) to determine telogen and anagen			
2=212	counts, or structural hair shaft abnormality			
97010	Application of a modality to 1 or more areas; hot or cold packs			
97012	Application of a modality to 1 or more areas; traction, mechanical			
97014	Application of a modality to 1 or more areas; electrical stimulation			
07017	(unattended)			
97016	Application of a modality to 1 or more areas; vasopneumatic devices			
97018	Application of a modality to 1 or more areas; paraffin bath			
97022	Application of a modality to 1 or more areas; whirlpool			
97024	Application of a modality to 1 or more areas; diathermy (eg, microwave)			
97026	Application of a modality to 1 or more areas; infrared			
97028	Application of a modality to 1 or more areas; ultraviolet			
97032	Application of a modality to 1 or more areas; electrical stimulation			
97033	(manual), each 15 minutes  Application of a modality to 1 or more areas; iontophoresis, each 15 minutes			
97033	Application of a modality to 1 or more areas; contrast baths, each 15 minutes  Application of a modality to 1 or more areas; contrast baths, each 15			
9703 <del>4</del>	minutes			
97035				
97036	Application of a modality to 1 or more areas; ultrasound, each 15 minutes  Application of a modality to 1 or more areas; Hubbard tank, each 15 minutes			
97129	Therapeutic interventions that focus on cognitive function (eg. attention,			
9/129	memory, reasoning, executive function, problem solving, and/or pragmatic			
	functioning) and compensatory strategies to manage the performance of an			
	activity (eg, managing time or schedules, initiating, organizing, and			
	sequencing tasks), direct (one-on-one) patient contact; initial 15 minutes			
97130	Therapeutic interventions that focus on cognitive function (eg, attention,			
,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	memory, reasoning, executive function, problem solving, and/or pragmatic			
	functioning) and compensatory strategies to manage the performance of an			
	Tanted onling   und compensation   Strategies to manage the performance of an			



<b>CPT</b> ®	Description
Codes	
	activity (eg, managing time or schedules, initiating, organizing, and
	sequencing tasks), direct (one-on-one) patient contact; each additional 15
	minutes (List separately in addition to code for primary procedure)
97151	Behavior identification assessment, administered by a physician or other
	qualified health care professional, each 15 minutes of the physician's or
	other qualified health care professional's time face-to-face with patient
	and/or guardian(s)/caregiver(s) administering assessments and discussing
	findings and recommendations, and non-face-to-face analyzing past data,
07150	scoring/interpreting the assessment, and preparing the report/treatment plan
97152	Behavior identification-supporting assessment, administered by one
	technician under the direction of a physician or other qualified health care
07152	professional, face-to-face with the patient, each 15 minutes
97153	Adaptive behavior treatment by protocol, administered by technician under
	the direction of a physician or other qualified health care professional, face- to-face with one patient, each 15 minutes
97154	Group adaptive behavior treatment by protocol, administered by technician
7/134	under the direction of a physician or other qualified health care professional,
	face-to-face with two or more patients, each 15 minutes
97155	Adaptive behavior treatment with protocol modification, administered by
7,100	physician or other qualified health care professional, which may include
	simultaneous direction of technician, face-to-face with one patient, each 15
	minutes
97156	Family adaptive behavior treatment guidance, administered by physician or
	other qualified health care professional (with or without the patient present),
	face-to-face with guardian(s)/caregiver(s), each 15 minutes
97157	Multiple-family group adaptive behavior treatment guidance, administered
	by physician or other qualified health care professional (without the patient
	present), face-to-face with multiple sets of guardians/caregivers, each 15
0=1.50	minutes
97158	Group adaptive behavior treatment with protocol modification, administered
	by physician or other qualified health care professional, face-to-face with
07520	multiple patients, each 15 minutes  Theremoving activities direct (one on one) retient contact (was of dynamic
97530	Therapeutic activities, direct (one-on-one) patient contact (use of dynamic activities to improve functional performance), each 15 minutes
07.522	1 //
97533	Sensory integrative techniques to enhance sensory processing and promote
	adaptive responses to environmental demands, direct (one-on-one) patient
97810	contact, each 15 minutes  Acupuncture, one or more needles, w/o electric stimulation; initial
9/010	15 minutes of personal one-one contact with the patient
97811	Acupuncture, one or more needles, w/o electric stimulation; each additional
7/011	15 minutes of personal one-one contact with the patient, with re-insertion of
	needles (s)
97813	Acupuncture, one or more needles, with electric stimulation; initial
<del></del>	15 minutes of personal one-one contact with the patient



CPT®	Description
Codes	
97814	Acupuncture, one or more needles, with electric stimulation; each additional
	15 minutes of personal one-one contact with the patient, with re-insertion of
	the needle(s) (List separately in addition to code for primary procedure)
98940	Chiropractic manipulative treatment (CMT); spinal, 1 to 2 regions
98941	Chiropractic manipulative treatment (CMT); spinal, 3 to 4 regions
98942	Chiropractic manipulative treatment (CMT); spinal, 5 regions
98943	Chiropractic manipulative treatment (CMT); extraspinal, 1 or more
	regions

### HCPCS codes considered not medically necessary when billed with a sole diagnosis of ADHD

HCPCS Codes	Description
G0176	Activity therapy, such as music, dance, art or play therapies not for recreation, related to the care and treatment of patient's disabling mental health problems, per session (45 minutes or more)
P2031	Hair analysis (excluding arsenic)
S8040	Topographic brain mapping

**ICD-10-CM Diagnosis Codes that Support Medical Necessity** 

ICD-10-CM Code	Description
F90.0 through F90.9	Attention-deficit hyperactivity disorders

Reviews, Revisions, and Approvals	Revision Date	Approval Date	Effective Date
Rebranded from corporate policy	5/23	9/13/23	10/13/23
Annual review. Added examples to II.A.5. under "Computerized electroencephalogram". Updated II.B.9 to "Cannabinoids and cannabinoid products". Removed "Supportive counseling" and "Vayarin" from II.B. Replaced "Endeavor RX" with "Video gamebased interventions (e.g., EndeavorRX, AKL-T01) in II.B.39. Added CPT codes 70544, 70545, 70546, 95957 as well as CPT codes 97151 through 97158 for ABA services included under II.B.23 "Intensive behavioral intervention programs" to not medically necessary table. Updated and reorganized background with no clinical significance. Coding reviewed. References reviewed and updated. Removed deleted CPT 92569 from table of CPT codes considered not medically necessary when billed with a sole diagnosis of ADHD.	03/25	6/9/25	7/10/25



#### References

- 1. Post RE, Kurlansik SL. Diagnosis and management of adult attention-deficit/hyperactivity disorder. *Am Fam Physician*. 2012;85(9):890 to 896.
- 2. Bukstein O. Attention deficit hyperactivity disorder in adults: Epidemiology, pathogenesis, clinical features, course, assessment, and diagnosis. UpToDate. <a href="www.uptodate.com">www.uptodate.com</a>. Updated November 27, 2023. Accessed January 4, 2024.
- 3. Krull KR. Attention deficit hyperactivity disorder in children and adolescents: Epidemiology and pathogenesis. UpToDate. <a href="https://www.uptodate.com">www.uptodate.com</a>. Updated March 9, 2023. Accessed January 4, 2024.
- Krull KR. Attention deficit hyperactivity disorder in children and adolescents: Overview of treatment and prognosis. UpToDate. <a href="www.uptodate.com">www.uptodate.com</a>. Updated September 19, 2023. Accessed January 4, 2024.
- 5. Wolraich ML, Hagan JF Jr, Allan C, et al. Clinical practice guideline for the diagnosis, evaluation, and treatment of attention-deficit/hyperactivity disorder in children and adolescents [published correction appears in Pediatrics. 2020 Mar;145(3):]. *Pediatrics*. 2019;144(4):e20192528. doi:10.1542/peds.2019 to 2528
- 6. Gibbins C, Weiss M. Clinical recommendations in current practice guidelines for diagnosis and treatment of ADHD in adults. *Curr Psychiatry Rep.* 2007;9(5):420 to 426. doi:10.1007/s11920-007-0055-1
- 7. Chan E, Fogler JM, Hammerness PG. Treatment of Attention-Deficit/Hyperactivity Disorder in Adolescents: A Systematic Review. *JAMA*. 2016;315(18):1997 to 2008. doi:10.1001/jama.2016.5453
- 8. Pliszka S; AACAP Work Group on Quality Issues. Practice parameter for the assessment and treatment of children and adolescents with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry*. 2007;46(7):894 to 921. doi:10.1097/chi.0b013e318054e724
- 9. Gloss D, Varma JK, Pringsheim T, Nuwer MR. Practice advisory: The utility of EEG theta/beta power ratio in ADHD diagnosis: Report of the guideline development, dissemination, and implementation subcommittee of the american academy of neurology. *Neurology*. 2016;87(22):2375 to 2379. doi:10.1212/WNL.000000000003265
- 10. Tseng PT, Cheng YS, Yen CF, et al. Peripheral iron levels in children with attention-deficit hyperactivity disorder: a systematic review and meta-analysis. *Sci Rep.* 2018;8(1):788. Published 2018 Jan 15. doi:10.1038/s41598-017-19096-x
- 11. Wang Y, Huang L, Zhang L, Qu Y, Mu D. Iron Status in Attention-Deficit/Hyperactivity Disorder: A Systematic Review and Meta-Analysis. *PLoS One*. 2017;12(1):e0169145. Published 2017 Jan 3. doi:10.1371/journal.pone.0169145
- 12. Krull KR. Attention deficit hyperactivity disorder in children and adolescents: Clinical features and diagnosis. UpToDate. <a href="www.uptodate.com">www.uptodate.com</a>. Updated March 9, 2023. Accessed January 4, 2024.
- 13. Dalrymple RA, McKenna Maxwell L, Russell S, Duthie J. NICE guideline review: Attention deficit hyperactivity disorder: diagnosis and management (NG87). *Arch Dis Child Educ Pract Ed.* 2020;105(5):289 to 293. doi:10.1136/archdischild-2019-316928
- 14. Barbaresi WJ, Campbell L, Diekroger EA, et al. Society for Developmental and Behavioral Pediatrics Clinical Practice Guideline for the Assessment and Treatment of Children and Adolescents with Complex Attention-Deficit/Hyperactivity Disorder. *J Dev Behav Pediatr*. 2020;41 Suppl 2S:S35 to S57. doi:10.1097/DBP.00000000000000770



- 15. Berger S. Cardiac evaluation of patients receiving pharmacotherapy for attention deficit hyperactivity disorder. UpToDate. <a href="www.uptodate.com">www.uptodate.com</a>. Updated March 16, 2023. Accessed January 4, 2024.
- 16. Not just ADHD? Helping children with multiple concerns. Centers for Disease Control and Prevention. Updated September 27, 2023. Accessed February 19, 2024.
- 17. Cândido RCF, Menezes de Padua CA, Golder S, Junqueira DR. Immediate-release methylphenidate for attention deficit hyperactivity disorder (ADHD) in adults. *Cochrane Database Syst Rev.* 2021;1(1):CD013011. Published 2021 Jan 18. doi:10.1002/14651858.CD013011.pub2
- 18. Brent D, Bukstein O, Solanto MV. Attention deficit hyperactivity disorder in adults: Treatment overview. UpToDate. <a href="www.uptodate.com">www.uptodate.com</a>. Updated August 10, 2023. Accessed January 4, 2024.
- 19. Young S, Hollingdale J, Absoud M, et al. Guidance for identification and treatment of individuals with attention deficit/hyperactivity disorder and autism spectrum disorder based upon expert consensus. *BMC Med.* 2020;18(1):146. doi:10.1186/s12916-020-01585-y
- 20. National Institute for Mental Health. Attention Deficit Hyperactivity Disorder. <a href="https://www.nimh.nih.gov/health/topics/attention-deficit-hyperactivity-disorder-adhd">https://www.nimh.nih.gov/health/topics/attention-deficit-hyperactivity-disorder-adhd</a>. Updated September 2023. Accessed January 4, 2024.

#### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.



This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care and are solely responsible for the medical advice and treatment of members/enrollees. This clinical policy is not intended to recommend treatment for members/enrollees. Members/enrollees should consult with their treating physician in regard to diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members/enrollees and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members/enrollees and their representatives agree to be bound by such terms and conditions by providing services to members/enrollees and/or submitting claims for payment for such services.

© 2023 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein.