

CONCERT GENETIC TESTING: EYE DISORDERS

Reference Number: LA.CP.CG.05 Date of Last Revision 12/23 Coding implications Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

OVERVIEW

In the past 15 years, genetics experts have identified approximately 500 genes that contribute to inherited eye diseases. Approximately 4,000 diseases affect humans, and nearly one-third of these diseases may affect the eyes. Because many genes involved in ophthalmologic disorders are now identified, scientists have developed a better understanding of how these genes influence vision and eye health.

Age-related macular degeneration (AMD) is an eye condition that causes damage to the central portion of the retina (the macula), affecting the ability to see objects straight ahead. It is a complex disease and is the leading cause of blindness and irreversible vision loss among adults over the age of 65 years. The etiology of AMD is multifactorial and includes both genetic and environmental (e.g., age, smoking) factors. Genetic testing has been proposed to predict the risk of developing advanced AMD in asymptomatic individuals, however, the clinical utility of genetic testing for age-related macular degeneration is limited. No studies have shown improvements in patients identified as being high-risk based on genetic testing, and evidence is insufficient to determine the effects of genetic testing on health outcomes. For individuals who have age-related macular degeneration, the clinical utility of genetic testing is limited and has not shown to be superior to clinical evaluation.

The molecular genetic basis for glaucoma has not been clearly elucidated, however a small subset of genes have been identified in very rare forms of congenital glaucoma.

Inherited retinal dystrophy can be caused by biallelic variants in the *RPE65* gene and other genes and can result in difficulty seeing in dim light and progressive loss of vision. Historically considered untreatable, gene therapy has been proposed as a treatment to improve visual function. Individuals who have vision loss due to biallelic *RPE65* variant associated retinal dystrophy are eligible to receive gene therapy. Because this is a rare condition, there are challenges with generating evidence demonstrating that the technology results in a meaningful improvement in net health outcomes.



POLICY REFERENCE TABLE

Coding Implications

This clinical policy references Current Procedural Terminology (CPT[®]). CPT[®] is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2022, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only and may not support medical necessity. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

NOTE: Coverage is subject to each requested code's inclusion on the corresponding LDH fee schedule. Non-covered codes are denoted (*) and are reviewed for Medical Necessity for members under 21 years of age on a per case basis.

Criteria Sections	Example Tests (Labs)	Common CPT Codes	Common ICD Codes	<u>Ref</u>				
Known Familial Variant Analysis for Eye Disorders								
<u>Known Familial</u> Variant Analysis for Eye Disorders	Targeted Mutation Analysis for a Known Familial Variant	81403*		6				
Macular Degeneration								
<u>Macular</u> Degeneration	Macula Risk (Arctic Medical Laboratories	81479, 81599*	H35.30, H35.3110-	2				
	Vita Risk (Arctic Medical Laboratories)	0205U*	H35.3194, H35.3210-					
	Macular Degeneration NGS Panel (Fulgent Genetics)	81404*, 81406*, 81408*, 81479	H35.3293, Z13.5					
RPE-Associated Re	etinal Dystrophy/Leber Congenital Amai	irosis						
and/or Deletion/Duplicatio n Analysis or	<i>RPE65</i> -Association Disorders via the <i>RPE65</i> Gene (PreventionGenetics, part of Exact Sciences)		H35.50- H35.54	1, 3, 4				
<u>Multigene Panel</u> <u>Analysis</u>	Leber Congenital Amaurosis Panel (PreventionGenetics, part of Exact Sciences)	81404*, 81406*, 81408*, 81479						

Please see the Concert Genetics Platform for a comprehensive list of registered tests.



<u>Glaucoma</u>						
<u>Glaucoma</u>	Glaucoma Panel (PreventionGenetics, part of Exact Sciences) Glaucoma Panel (Blueprint Genetics)	81404*, 81406*, 81407*, 81408*, 81479		1		
Other Covered Eye Disorders						
Other Covered Eye Disorders	See below	81400*-81408*		1, 5, 7, 8		

OTHER RELATED POLICIES

This policy document provides criteria for Genetic Testing for Eye Disorders. Please refer to:

- *Genetic Testing: Hereditary Cancer Susceptibility* for criteria related to genetic testing for retinoblastoma.
- *Genetic Testing: Hearing Loss* for criteria related to genetic testing for disorders that include hearing loss, such as Usher syndrome.
- *Genetic Testing: Multisystem Inherited Disorders, Intellectual Disability, and Developmental Delay* for criteria related to oculocutaneous albinism and other multisystem inherited disorders.
- *Genetic Testing: General Approach to Genetic and Molecular Testing* for criteria related to genetic testing for eye disorders that are not specifically discussed in this or another non-general policy.

CRITERIA

It is the policy of Louisiana Healthcare Connections that the specific genetic testing noted below is **medically necessary** when meeting the related criteria:

KNOWN FAMILIAL VARIANT ANALYSIS FOR EYE DISORDERS

- I. Targeted mutation analysis for a known familial variant (81403*) for an eye disorder is considered **medically necessary** when:
 - A. The member/enrollee has a <u>close relative</u> with a known pathogenic or likely pathogenic variant causing the condition.



II. Targeted mutation analysis for a known familial variant (81403*) for an eye disorder is considered **investigational** for all other indications.

back to top

MACULAR DEGENERATION

I. Genetic testing for <u>macular degeneration</u> (81404*, 81406*, 81408*, 81479, 81599*, 0205U*) is considered **investigational**.

back to top

RPE65-ASSOCIATED RETINAL DYSTROPHY / LEBER CONGENITAL AMAUROSIS

RPE65 Sequencing and/or Deletion/Duplication Analysis or Multigene Panel Analysis

- Genetic testing for <u>RPE65-associated retinal dystrophy/Leber congenital amaurosis</u> via RPE65 sequencing and/or deletion/duplication analysis (81406*, 81479) or a multigene panel (81404*, 81406*, 81408*, 81479) that includes RPE65 is considered **medically** necessary when:
 - A. The member/enrollee has a diagnosis of a retinal dystrophy or Leber Congenital Amaurosis, **AND**
 - B. The member/enrollee is being considered for treatment with voretigene neparvovec (Luxturna®).
- II. Genetic testing for <u>RPE65-associated retinal dystrophy/Leber congenital amaurosis</u> via *RPE65* sequencing and/or deletion/duplication analysis (81406*, 81479) or a multigene panel (81404*, 81406*, 81408*, 81479) that includes *RPE65* is considered investigational for all other indications.

back to top

GLAUCOMA

I. Genetic testing for glaucoma (81404*, 81406*, 81407*, 81408*, 81479) is considered investigational.

back to top



OTHER COVERED EYE DISORDERS

Genetic Testing Guidelines for Genetic Eye Disorders

The following is a list of conditions that have a known genetic association. Due to their relative rareness, it may be appropriate to approve claims for these tests to establish or confirm a diagnosis.

- I. Genetic testing to establish or confirm one of the following eye disorders to guide management is considered **medically necessary** when the member/enrollee demonstrates clinical features** consistent with the disorder (the list is not meant to be comprehensive, see II below):
 - A. Duane Syndrome
 - B. Familial Exudative Vitreoretinopathy
 - C. <u>Retinitis Pigmentosa</u>
 - D. <u>Aniridia</u>
 - E. X-linked Congenital Retinoschisis
 - F. Presenile Cataracts
- II. Genetic testing to establish or confirm the diagnosis of all other eye disorders not specifically discussed within this or another medical policy will be evaluated by the criteria outlined in *General Approach to Genetic and Molecular Testing* (see policy for criteria).

**Clinical features for a specific disorder may be outlined in resources such as <u>GeneReviews</u>, <u>OMIM</u>, <u>National Library</u> <u>of Medicine</u>, <u>Genetics Home Reference</u>, or other scholarly sources.

back to top

NOTES AND DEFINITIONS

- 1. Close relatives include first, second, and third degree blood relatives on the same side of the family:
 - a. First-degree relatives are parents, siblings, and children
 - b. **Second-degree relatives** are grandparents, aunts, uncles, nieces, nephews, grandchildren, and half siblings
 - c. **Third-degree relatives** are great grandparents, great aunts, great uncles, great grandchildren, and first cousins
- 2. Age-related Macular Degeneration (AMD) is the leading cause of blindness and irreversible vision loss among older adults (greater than age 65 years).



- 3. **Retinal dystrophies (RDs)** are degenerative diseases of the retina which have marked clinical and genetic heterogeneity. Vision impairment may vary from poor peripheral or night vision to complete blindness, and severity usually increases with age.
- 4. *RPE65* (retinal pigment epithelium-specific protein 65-kD) gene encodes the RPE54 protein, which is an all translate-retinal isomerase, a key enzyme expressed in the retinal pigment epithelium (RPE) that is responsible for regeneration of 11-cis-retinol in the visual cycle.
- 5. **Gene therapy** are treatments that change the expression of genes to treat disease, e.g., by replacing or inactivating a gene that is not functioning properly or by introducing a new gene. Genes may be introduced into human cells through a vector, usually a virus.

back to top

CLINICAL CONSIDERATIONS

The purpose of genetic testing of asymptomatic individuals with risk of developing age-related macular degeneration is to identify single nucleotide variants for primary prevention or earlier detection of disease for more timely intervention to affect course of disease progression. Patients may be referred from primary care to an ophthalmologist or medical geneticist for investigation and management of age-related macular degeneration. In all cases, the patient should receive counseling from a physician with expertise in inherited disease or a genetic counselor. Whenever clinical findings suggest the presence of an inherited eye disease, the treating ophthalmologist should either discuss the potential value of genetic testing with their patient and order the appropriate tests (if any) or should offer a referral to another physician or counselor with expertise in the selection and interpretation of genetic tests. Treating physicians should also ensure that their patients receive a written copy of their genetic test results.

Genetic testing is required to detect the presence of pathogenic or likely pathogenic variants in the *RPE65* gene in individuals with documented vision loss. By definition, pathogenic or likely pathogenic variant(s) must be present in both copies of the *RPE65* gene to establish a diagnosis of biallelic *RPE65*-mediated inherited retinal dystrophy. Next-generation sequencing and Sanger sequencing typically cannot resolve the phase (e.g., *trans* vs. *cis* configuration) when two *RPE65* pathogenic or likely pathogenic variants are detected. In this scenario, additional documentation of the *trans* configuration is required to establish a diagnosis of biallelic *RPE65*-mediated inherited retinal dystrophy.

back to top

BACKGROUND AND RATIONALE

Known Familial Variant Analysis for Eye Disorders

Genetic Support Foundation



The Genetic Support Foundation's Genetics 101 information on genetic testing says the following about testing for familial pathogenic variants:

Genetic testing for someone who may be at risk for an inherited disease is always easier if we know the specific genetic cause. Oftentimes, the best way to find the genetic cause is to start by testing someone in the family who is known or strongly suspected to have the disease. If their testing is positive, then we can say that we have found the familial pathogenic (harmful) variant. We can use this as a marker to test other members of the family to see who is also at risk.

Macular Degeneration

American Society of Retina Specialists

American Society of Retina Specialists (2017) published special correspondence on the use of genetic testing in the management of patients with age-related macular degeneration, which made the following conclusions:

- 1. Age-related macular degeneration (AMD) genetic testing may provide information on the progression rates from intermediate to advanced AMD. However, before ordering this testing, retina specialists should be aware of the following:
 - a. Testing should be performed only at Clinical Laboratory Improvement Amendments-certified laboratories with expertise in genetic sequencing. Because of the high variability in the results, direct-to-consumer (DTC) AMD genetic testing that does not meet this standard is not recommended.
 - b. Interpretation of the results of AMD genetic testing is complex.
 - c. At present, there is no clinical evidence that altering the management of genetically higher risk progression patients, for example, with more frequent office visits and/or improved lifestyle changes, results in better visual outcomes for these patients compared with individuals of lower genetic susceptibility. As such, prospective studies are needed before patient care is modified.
- 2. Age-related macular degeneration genetic testing at present in patients with neovascular AMD does not provide clinically relevant information regarding response to anti-vascular endothelial growth factor (VEGF) treatment and is not recommended for this purpose.
- **3**. Although genetic testing to determine the optimal nutritional supplementation may in the future prove useful, at present there is insufficient data to support the use of genetic testing in patients with AMD prior to recommendation of current Age-Related Eye Disease Study (AREDS) nutritional supplement use. (p. 75)

RPE65 Sequencing and/or Deletion/Duplication Analysis or Multigene Panel Analysis

US Food and Drug Administration (FDA)



The FDA issued an approval letter on December 18, 2017, for Luxturna stating, "Under this license, you are authorized to manufacture the product voretigene neparvovec-rzyl, which is indicated for the treatment of patients with confirmed biallelic *RPE65* mutation-associated retinal dystrophy." (p. 1)

National Institute for Health and Care Excellence

The National Institute for Health and Care Excellence published guidance for the use of voretigene neparvovec for treating inherited retinal dystrophies caused by *RPE65* gene mutations, which stated the following:

1.1 Voretigene neparvovec is recommended, within its marketing authorisation, as an option for treating *RPE65*-mediated inherited retinal dystrophies in people with vision loss caused by inherited retinal dystrophy from confirmed biallelic *RPE65* mutations and who have sufficient viable retinal cells. It is recommended only if the company provides voretigene neparvovec according to the commercial arrangement. (p. 4)

The committee noted that, "*RPE65*-mediated inherited retinal dystrophies are rare and serious. They involve progressive loss of vision. This ultimately leads to near-total blindness, and severely affects the quality of life of people with the condition, and their families and carers. Current treatment is supportive care. Clinical trial evidence shows that, in the short term, voretigene neparvovec improves vision and prevents the condition from getting worse." (p. 4)

American Academy of Ophthalmology (AAO)

The American Academy of Ophthalmology (AAO) Task Force on Genetic Testing published recommendations for genetic testing of inherited eye diseases (2014). In it, they state "there are some situations in which limited parallel testing is the most effective strategy. When a clinical disease is caused by multiple different genes (e.g., nonsyndromic retinitis pigmentosa, Usher syndrome, Leber congenital amaurosis, and Bardet Biedl syndrome), it often is best to order a single test that has been designed specifically to evaluate efficiently all of the genes known to cause the patient's clinical findings". (p. 3) They also recommend that one should "offer genetic testing to patients with clinical findings suggestive of a Mendelian disorder whose causative gene(s) have been identified." (p. 4)

Glaucoma

American Academy of Ophthalmology (AAO)

The American Academy of Ophthalmology (AAO) Task Force on Genetic Testing published recommendations for genetic testing of inherited eye diseases (2014) which stated, in part: "Avoid routine genetic testing for genetically complex disorders like age-related macular degeneration and late-onset primary open-angle glaucoma until specific treatment or surveillance strategies have been shown in 1 or more published prospective clinical trials to be of benefit to individuals with specific disease-associated genotypes. In the meantime, confine the genotyping of such patients to research studies." (p. 5)



OTHER COVERED EYE DISORDERS

General Testing Guidelines for Genetic Eye Disorders

American Academy of Ophthalmology (AAO)

The American Academy of Ophthalmology (AAO) Task Force on Genetic Testing published the following recommendations for genetic testing of inherited eye diseases (2014):

- 1. Offer genetic testing to patients with clinical findings suggestive of a Mendelian disorder whose causative gene(s) have been identified. If unfamiliar with such testing, refer the patient to a physician or counselor who is. In all cases, ensure that the patient receives counseling from a physician with expertise in inherited disease or a certified genetic counselor.
- 2. Use Clinical Laboratories Improvement Amendments– approved laboratories for all clinical testing. When possible, use laboratories that include in their reports estimates of the pathogenicity of observed genetic variants that are based on a review of the medical literature and databases of disease-causing and non–disease-causing variants.
- 3. Provide a copy of each genetic test report to the patient so that she or he will be able independently to seek mechanism-specific information, such as the availability of gene-specific clinical trials, should the patient wish to do so.
- 4. Avoid direct-to-consumer genetic testing and discourage patients from obtaining such tests themselves. Encourage the involvement of a trained physician, genetic counselor, or both for all genetic tests so that appropriate interpretation and counseling can be provided.
- 5. Avoid unnecessary parallel testing— order the most specific test(s) available given the patient's clinical findings. Restrict massively parallel strategies like whole-exome sequencing and whole-genome sequencing to research studies conducted at tertiary care facilities.
- 6. Avoid routine genetic testing for genetically complex disorders like age-related macular degeneration and late-onset primary open-angle glaucoma until specific treatment or surveillance strategies have been shown in 1 or more published prospective clinical trials to be of benefit to individuals with specific disease-associated genotypes. In the meantime, confine the genotyping of such patients to research studies.
- 7. Avoid testing asymptomatic minors for untreatable disorders except in extraordinary circumstances. For the few cases in which such testing is believed to be warranted, the following steps should be taken before the test is performed: (1) the parents and child should undergo formal genetic counseling, (2) the certified counselor or physician performing the counseling should state his or her opinion in writing that the test is in the family's best interest, and (3) all parents with custodial responsibility for the child should agree in writing with the decision to perform the test. (p. 4 and 5)

back to top



	connections	
Reviews, Revisions, and Approvals	Revision Date	Approval Date
Policy developed.	09/23	11/27/23
Semi-annual review. Overview, coding, reference-table, background and references updated. Throughout policy: replaced "coverage criteria" with "criteria. For Policy Reference Table: removed "Stargardt Disease" and added "Macular Degeneration"; under RPE-Associated Retinal Dystrophy/Leber Congenital Amaurosis and Glaucoma: added "part of Exact Sciences"; added "Other" to Covered Eye Disorders. For Other Related Policies: added "Molecular" to Genetic Testing: General Approach to Genetic Testing. For RPE65-Associated Retinal Dystrophy/Leber Congenital Amaurosis: in II. added "81479". For Notes and Definitions: 1. added "on the same side of the family". For Clinical Considerations: removed "In all cases, the patient should receive". For Background and Rationale: removed "inheritance patterns" and replaced with "genetic testing".	12/23	2/27/24

REFERENCES

- American Academy of Ophthalmology Task Force on Genetic Testing. Recommendations for Genetic Testing of Inherited Eye Diseases - 2014. Accessed 11/11/2021. <u>https://www.aao.org/clinical-statement/recommendations-genetic-testing-of-inherited-eye-d</u>
- Csaky KG SA, Kaiser PK, et al. The Use of Genetic Testing in the Management of Patients with Age-Related Macular Degeneration: American Society of Retina Specialists Genetics Task Force Special Report. 2017. Accessed November 2, 2020.
- 3. US Food & Drug Administration. Approval Letter LUXTURNA. December 19, 2017. Accessed November 2, 2020. <u>https://www.fda.gov/media/109487/download</u>
- National Institute for Health and Care Excellence (NICE). Voretigene neparvovec for treating inherited retinal dystrophies caused by RPE65 gene mutations [HST11]. October 2019. Accessed November 2, 2020. <u>http://nice.org.uk/guidance/hst11</u>
- 5. MedlinePlus [Internet]. Bethesda (MD): National Library of Medicine (US). Available from: <u>https://medlineplus.gov/genetics/</u>
- 6. Genetic Support Foundation. Genetics 101 Genetic Testing: Familial Pathogenic Variant. Accessed 10/4/2022. <u>https://geneticsupportfoundation.org/genetics-101/#</u>
- Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2023. Available from: <u>https://www.ncbi.nlm.nih.gov/books/NBK1116/</u>
- Online Mendelian Inheritance in Man, OMIM. McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University (Baltimore, MD). World Wide Web URL: <u>https://omim.org/</u>

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program



approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members/enrollees. This clinical policy is not intended to recommend treatment for members/enrollees. Members/enrollees should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members/enrollees and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members/enrollees and their representatives agree to be bound by such terms and conditions by providing services to members/enrollees and/or submitting claims for payment for such services.

©2023 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained



herein. Louisiana Healthcare Connections is a registered trademarks exclusively owned by Louisiana Healthcare Connections.

back to top