

# Clinical Policy: Bronchial Thermoplasty

Reference Number: LA.CP.MP.110

Date of Last Revision: 11/2021

Coding Implications  
Revision Log

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

## Description

This policy describes the medical necessity requirements for bronchial thermoplasty (BT). BT is a bronchoscopic procedure that utilizes radiofrequency ablation to reduce airway smooth muscle cells. It is designed to serve as a therapeutic option to reduce severe bronchoconstriction for severe persistent asthma.

## Policy/Criteria

It is the policy of Louisiana Healthcare Connections that the long-term safety and effectiveness of bronchial thermoplasty has not been proven for severe asthma or any other indications.

## Background

Asthma is a common inflammatory syndrome caused by chronic, intermittent obstruction of the lower respiratory tract that affects millions of individuals. This process is mediated by several inflammatory cytokines, chemokines, adhesion molecules, and signal transduction cascades.<sup>1</sup> T helper type 2 (T<sub>H</sub>2) and type 17 (T<sub>H</sub>17) CD4<sup>+</sup>, basophils, eosinophils, mast cells, and type 2 innate lymphoid cells are crucial for mediating the asthmatic response.<sup>2</sup>

BT is a bronchoscopic procedure that applies thermal energy to the airway wall and, thereby, reduces the extent of airway smooth muscle cell hypertrophy via radiofrequency ablation.<sup>3</sup> Some studies published on BT have tested its therapeutic potential against severe asthma.<sup>4</sup> However, the literature recently published on BT has been controversial and the studies evaluating the efficacy of BT have not provided consistent results.

A prospective non-randomized study of 16 patients with stable mild to moderate asthma found a significant reduction in airway hyperresponsiveness without a change in FEV<sub>1</sub>.<sup>5</sup> The Asthma Intervention Research Trial (AIR), a randomized controlled trial that enrolled 112 patients, showed an improvement in asthma symptoms from BT but no reduction in hyperresponsiveness or FEV<sub>1</sub>.<sup>6</sup> The Research in Severe Asthma Trial (RISA), a small randomized study that enrolled only 32 patients, assessed the safety of BT in patients receiving high doses of steroids. Despite several complications, including hospitalizations, a difference was seen in the BT group versus control.<sup>7</sup> Some critics argue that these studies lack the statistical power and blinded placebo control to demonstrate clear conclusions on the efficacy of BT's clinical potential.<sup>8</sup>

In 2010, Castro *et al.* performed a randomized, controlled trial with 288 patients that included a placebo control. This study was called the Asthma Intervention Research Trial 2 (AIR2).<sup>9</sup> AIR2 found a statistically significant improvement in their primary outcome, which was the score from the Asthma Quality of Life Questionnaire (AQLQ).<sup>9</sup> However, these scores fell below a clinically meaningful threshold.<sup>4</sup> There was no difference in peak flow, rescue medication use, or FEV<sub>1</sub>.<sup>9</sup> Moreover, several investigators have criticized the AIR2 study for failing to meet secondary outcome measures such as safety, its patient selection, and its true efficacy.<sup>8,10,11</sup> Thus, this study also remains controversial.

Hayes conducted a review of the available literature on BT, noting that overall the body of evidence is small and of low quality. The findings were that BT may improve quality of life outcomes, however other results including emergency department visits, symptom relief, and medication use were inconsistent across studies. BT was found to not decrease hospitalization following treatment and it actually increased hospitalization during the treatment period. Treatment with BT was associated with a statistically significant increase in complications such as wheezing, chest discomfort, night awakenings, sputum discoloration, and cough. The quality and quantity of evidence was not enough to establish the long-term safety and efficacy of the procedure.<sup>12</sup>

A meta-analysis of the aforementioned randomized, controlled trials by Wu, et al, suggests that while BT significantly improves AQLQ scores, there were more respiratory adverse events and hospitalizations for respiratory adverse events with BT than with medications or with placebo.<sup>13</sup>

Studies at 5 year follow up have reported BT to be safe (stable pulmonary function test and no bronchiectasis on chest CT) with persistent reductions in asthma exacerbation rates and/or emergency department visits/hospitalizations.<sup>21,22</sup> The complexity and uncertainties in the selection of patients for BT require a multidisciplinary team approach at asthma centers with high volumes of severe asthma patients and a high level of experience in interventional pulmonology procedures.<sup>21</sup>

#### ***European Respiratory Society/American Thoracic Society***

A 2014 joint statement by the European Respiratory Society and American Thoracic Society strongly recommends that BT be performed only in adults with severe asthma, in the context of a clinical trial or independent systematic registry. They conclude that the body of evidence is of very low quality, and that long-term benefits and safety are unknown.

#### ***National Institute for Health and Care Excellence (NICE)***

NICE guidance states that current evidence on the safety and efficacy of BT for severe asthma is adequate to support the use of this procedure provided that standard arrangements are in place for clinical governance, consent and audit. BT should only be done by clinicians with training in the procedure and experience in managing severe asthma. Further research should report details of patient selection and long-term safety and efficacy outcomes.

#### ***Global Initiative for Asthma***

The Global Initiative for Asthma recommends BT as a potential option for highly selected adult patients who have uncontrolled asthma despite use of recommended therapeutic regimens and referral to an asthma specialty center. Caution should be used in selecting patients for this procedure. BT should be performed in adults with severe asthma only in the context of an independent Institutional Review Board-approved systematic registry or a clinical study, so that further evidence about effectiveness and safety of the procedure can be accumulated. Longer-term follow-up of larger cohorts comparing effectiveness and safety, including lung function, in both active and sham- treated patients is needed.<sup>17</sup>

#### ***Agency for Healthcare Research and Quality***

Three RCTs and several descriptive studies meeting our inclusion criteria have evaluated BT. Based on the available literature, BT may be modestly beneficial in some patients with asthma, but is not without risks in any population. The risk of adverse events is higher during the treatment period and for several weeks afterward. Benefit is typically observed weeks to months after therapy and can last for at least 5 years, after which the duration of effect is unknown.

### ***British Thoracic Society***

Further research is needed to identify which patients with asthma might benefit from BT. However, it is likely that patients who remain uncontrolled despite optimal medical treatment and who have been considered for biological treatments and are either unsuitable for or fail a trial of such a treatment may be an appropriate group, as other treatment options for these patients are elusive. There are no trials comparing the efficacy of BT with biological treatments for people with asthma. BT may be considered for the treatment of adult patients (aged 18 and over) with severe asthma who have poorly-controlled asthma despite optimal medical therapy. An asthma specialist with expertise in BT should assess patients prior to undergoing treatment, and treatment should take place in a specialist centre with the appropriate resources and training, including access to an intensive care unit.<sup>20</sup>

### **Coding Implications**

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2020, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only and may not support medical necessity. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
31660	Bronchoscopy, rigid or flexible, including fluoroscopic guidance, when performed; with bronchial thermoplasty, 1 lobe
31661	Bronchoscopy, rigid or flexible, including fluoroscopic guidance, when performed; with bronchial thermoplasty, 2 or more lobes

Reviews, Revisions, and Approvals	Review Date	Approval Date
Converted corporate to local policy.	08/15/2020	
“Experimental/investigational” verbiage replaced in policy statement with descriptive language.” Removed ICD-10 table. Replaced all instances of “members” with “members/enrollees”. References reviewed and updated.	11/21	

## References

1. Barnes, Peter J. "Immunology of asthma and chronic obstructive pulmonary disease." *Nature Reviews Immunology* 8.3 (2008): 183-192.
2. Lambrecht, Bart N., and Hamida Hammad. "The immunology of asthma." *Nature immunology* 16.1 (2015): 45-56.
3. Miller, John D., et al. "A prospective feasibility study of bronchial thermoplasty in the human airway." *CHEST Journal* 127.6 (2005): 1999-2006.
4. Wahidi, Momen M., and Monica Kraft. "Bronchial thermoplasty for severe asthma." *American journal of respiratory and critical care medicine* 185.7 (2012): 709-714.
5. Cox, Gerard, et al. "Bronchial thermoplasty for asthma." *American journal of respiratory and critical care medicine* 173.9 (2006): 965-969.
6. Cox, Gerard, et al. "Asthma control during the year after bronchial thermoplasty." *New England journal of medicine* 356.13 (2007): 1327-1337.
7. Pavord, Ian D., et al. "Safety and efficacy of bronchial thermoplasty in symptomatic, severe asthma." *American journal of respiratory and critical care medicine* 176.12 (2007): 1185-1191.
8. Bel, Elisabeth H. "Bronchial thermoplasty: has the promise been met?." *American journal of respiratory and critical care medicine* 181.2 (2010): 101-102.
9. Castro, Mario, et al. "Effectiveness and safety of bronchial thermoplasty in the treatment of severe asthma: a multicenter, randomized, double-blind, sham-controlled clinical trial." *American journal of respiratory and critical care medicine* 181.2 (2010): 116-124.
10. Laxmanan, Balaji, et al. "Recent Advances in Chest Medicine: Advances in Bronchial Thermoplasty." *Chest* (2016).
11. Iyer, Vivek N., and Kaiser G. Lim. "Bronchial thermoplasty: Where there is smoke, there is fire." *Allergy and Asthma Proceedings*. Vol. 36. No. 4. OceanSide Publications, Inc, 2015.
12. Hayes Health Technology Assessment. Bronchial Thermoplasty for Treatment of Asthma. Annual Review June 19, 2019. Published May 26, 2016. Accessed March 29, 2021.
13. Wu, Q., et al. "Meta-analysis of the efficacy and safety of bronchial thermoplasty in patients with moderate-to-severe persistent asthma." *Journal of International Medical Research* 39.1 (2011): 10-22.
14. O'Reilly A, Browne I, Watchorn D, Egan JJ, Lane S. The efficacy and safety of bronchial thermoplasty in severe persistent asthma on extended follow-up. *QJM*. 2017 Nov 15. doi: 10.1093/qjmed/hcx221.
15. Chung KF, Wenzel SE, Brozek JL, et al. International ERS/ATS guidelines on definition, evaluation and treatment of severe asthma. *European respiratory journal*. 2014 43: 343-373; DOI: 10.1183/09031936.00202013
16. National Institute for Health and Care Excellence. Bronchial thermoplasty for severe asthma. Interventional procedures guidance IPG635. December 2018.
17. Global Initiative for Asthma (GINA). Global Strategy for Asthma Management and Prevention. Updated 2020.
18. Wenzel S. Treatment of severe asthma in adolescents and adults. In: UpToDate. Bochner BS (Ed). In: UpToDate, Waltham, MA. Accessed March 29, 2021
19. Agency for Healthcare Research and Quality. Effectiveness and Safety of Bronchial Thermoplasty in Management of Asthma. Comparative Effectiveness Review. Number 202. Dec 2017.

20. British Thoracic Society. SIGN 158. British guideline on the management of asthma. July 2019
21. Bonta PI, Chanez P, Annema JT, et al. Bronchial Thermoplasty in Severe Asthma: Best Practice Recommendations from an Expert Panel. *Respiration*. 2018;95(5):289-300.
22. Zhou JP, Feng Y, Wang Q, et al. Long-term efficacy and safety of bronchial thermoplasty in patients with moderate-to-severe persistent asthma: a systemic review and meta-analysis. *J Asthma*. 2016;53(1):94-100. doi: 10.3109/02770903.2015.1065424.

### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of member/enrollee. This clinical policy is not intended to recommend treatment for member/enrollee. Member/enrollee should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, member/enrollee and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, member/enrollee and their representatives agree to be bound by such terms and conditions by providing services to member/enrollee and/or submitting claims for payment for such services.

©2020 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.