

Clinical Policy: 25-hydroxyvitamin D Testing in Children and Adolescents

Reference Number: LA.CP.MP.157

Date of Last Review: 10/23

Coding Implications
Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

#### **Description**

A global consensus statement recommends against universal screening for vitamin D deficiency in healthy children as there is insufficient evidence that the potential benefits of testing outweigh the potential harms.<sup>2</sup>

#### Policy/Criteria

I. It is the policy of Louisiana Healthcare Connections that 25-hydroxyvitamin D testing in healthy, including obese but otherwise healthy, children (age  $\geq 1$  and  $\leq 18$ ) is **not medically necessary** because these tests have not been demonstrated to have a clear clinical benefit.

#### **Background**

Measurement of 25-OH-D (25-hydroxyvitamin D) concentration is the appropriate screening test for vitamin D deficiency. The 1,25-OH<sub>2</sub>-D test has little to no predictive value related to bone health.<sup>6</sup> However, there is lack of agreement concerning the best type of assay to conduct when measuring 25-hydroxyvitamin D.<sup>4</sup> Furthermore, there is substantial controversy concerning cutoff levels to define vitamin D deficiency, as the evidence is inconsistent regarding optimal levels of vitamin D.<sup>7,10</sup> The international Vitamin D Standardization Program has established measures for standardizing the laboratory value of 25(OH)D to improve clinical and public health practice.<sup>11</sup>

Prevalence of vitamin D deficiency in children (defined in the study as levels < 20 ng/mL) is approximately 15%, with estimates ranging from 14% to 37%. Rates of deficiency vary among certain populations, with increased risk among black and Hispanic teenagers, as well as overweight and obese children and adolescents. Reduced serum vitamin D in overweight and obese children and adolescents reflects sequestration in adipose tissue, but little is known about the significance of low serum vitamin D in this population.

A global consensus of 33 experts, convened at the request of the European Society for Pediatric Endocrinology, reviewed the available literature on prevention and management of nutritional rickets and determined that routine vitamin D screening is not recommended for healthy children. They note the frequent coexistence of dietary calcium and vitamin D deficiency, which alters the threshold for development of rickets and makes a single screening value impractical. The global consensus panel advocates for identification and screening of groups at high risk for vitamin D deficiency based on clinical factors, as opposed to universal screening of asymptomatic individuals as public health policy. 1,14

The American Academy of Pediatrics (AAP) – Section on Endocrinology advises against ordering vitamin D concentrations routinely in otherwise healthy children, including children who are overweight or obese.<sup>5</sup> Additionally, the Italian Pediatric Society and the Italian Society of Preventative and Social Pediatrics recommend against routine 25(OH) D testing in children and adolescents.<sup>13</sup> The AAP's report on optimizing bone health recommends screening for vitamin D



deficiency only in children and adolescents with conditions associated with reduced bone mass and/or recurrent low-impact fractures.<sup>6</sup> The Italian Pediatric Society and the Italian Society of Preventive and Social Pediatrics suggest measuring serum 25(OH) D levels in the presence of multiple risk factors for vitamin D deficiency.<sup>13</sup>

For healthy children and adolescents who are not ingesting enough foods with vitamin D, the Endocrine Society's clinical practice guidelines for the prevention of vitamin D deficiency and the AAP recommend supplementation with vitamin D, as does the global consensus panel convened by the European Society for Pediatric Endocrinology.<sup>2,6,7</sup>

### **Coding Implications**

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2022, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage and may not support medical necessity. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

Table 1: CPT codes not medically necessary when billed with a corresponding ICD-10-CM diagnosis code in Table 2.

<b>CPT</b> ®	Description
Codes	
82306	Vitamin D; 25 hydroxy, includes fraction(s), if performed

Table 2: ICD-10-CM diagnosis codes not medically necessary when billed with a corresponding CPT code in Table 1.

ICD-10-CM	Description	
Code		
E66.01	Morbid (severe) obesity due to excess calories	
E66.09	Other obesity due to excess calories	
E66.1	Drug-induced obesity	
E66.3	Overweight	
E66.8	Other obesity	
E66.9	Obesity, unspecified	
Z00.00	Encounter for general adult medical examination without abnormal findings	
Z00.129	Encounter for routine child health examination without abnormal findings	
Z00.8	Encounter for other general examination	
Z68.52	Body mass index (BMI) pediatric, 5 <sup>th</sup> percentile to less than 85 <sup>th</sup> percentile for	
	age	
Z68.53	BMI pediatric, 85 <sup>th</sup> percentile to less than 95 <sup>th</sup> percentile for age	
Z68.54	BMI pediatric, greater than or equal to 95 <sup>th</sup> percentile for age	



Reviews, Revisions, and Approvals	Revision Date	Approval
		Date
Converted corporate to local policy.	08/15/2020	
Annual review. Replaced "member" with "member/enrollee"	10/22	1/14/23
References reviewed and updated. Reviewed by specialist.		
Changed "Last Review Date" in the header to "Date of Last		
Review" and "Date" in revision log to "Revision Date". Updated		
background with no impact to criteria.		
Annual review completed. Background updated with no impact to	10/23	1/5/24
criteria. References reviewed and updated. External and internal		
specialist review.		

#### References

- 1. Munns CF, Shaw N, Kiely M, et al. Global Consensus Recommendations on Prevention and Management of Nutritional Rickets. *J Clin Endocrinol Metab*. 2016;101(2):394 to 415. doi:10.1210/jc.2015-2175
- 2. Saintonge S, Bang H, Gerber LM. Implications of a new definition of vitamin D deficiency in a multiracial us adolescent population: the National Health and Nutrition Examination Survey III. *Pediatrics*. 2009;123(3):797 to 803. doi:10.1542/peds.2008-1195
- 3. Misra M. Vitamin D insufficiency and deficiency in children and adolescents. UpToDate. <a href="https://www.uptodate.com">www.uptodate.com</a>. Published April 12, 2022. Accessed August 4, 2023.
- 4. Golden NH, Abrams SA; Committee on Nutrition. Optimizing bone health in children and adolescents. *Pediatrics*. 2014;134(4):e1229 to e1243. doi:10.1542/peds.2014-2173
- 5. Turer CB, Lin H, Flores G. Prevalence of vitamin D deficiency among overweight and obese US children. *Pediatrics*. 2013;131(1):e152 to e161. doi:10.1542/peds.2012-1711
- 6. Jin J. Screening for Vitamin D Deficiency in Adults. *JAMA*. 2021;325(14):1480. doi:10.1001/jama.2021.4606
- 7. Holick MF, Binkley NC, Bischoff-Ferrari HA, et al. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline [published correction appears in J Clin Endocrinol Metab. 2011 Dec;96(12):3908]. *J Clin Endocrinol Metab*. 2011;96(7):1911 to 1930. doi:10.1210/jc.2011-0385
- 8. Grossman Z, Hadjipanayis A, Stiris T, et al. Vitamin D in European children-statement from the European Academy of Paediatrics (EAP). *Eur J Pediatr*. 2017;176(6):829 to 831. doi:10.1007/s00431-017-2903-2
- 9. Canadian Association of Pathologists. Choosing Wisely Canada. Don't perform population based screening for 25-OH-Vitamin D deficiency. <a href="https://choosingwiselycanada.org/recommendation/pathology/">https://choosingwiselycanada.org/recommendation/pathology/</a>. Updated August 2022. Accessed August 4, 2023.
- Castano L, Madariaga L, Grau G, García-Castaño A. 25(OH)Vitamin D Deficiency and Calcifediol Treatment in Pediatrics. *Nutrients*. 2022;14(9):1854. Published 2022 Apr 29. doi:10.3390/nu14091854
- 11. Centers for Disease Control and Prevention (CDC). Vitamin D Standardization-Certification Program.
  - https://www.cdc.gov/labstandards/csp/vdscp.html#:~:text=What%20is%20the%20function%20of,a%20certain%20accuracy%20and%20precision. Updated March 9, 2023. Accessed August 4, 2023.



- 12. Loyal J, Cameron A. Vitamin D in Children: Can We Do Better? *Pediatrics*. 2020;145(6):e20200504. doi:10.1542/peds.2020-0504
- 13. Saggese G, Vierucci F, Prodam F, et al. Vitamin D in pediatric age: consensus of the Italian Pediatric Society and the Italian Society of Preventive and Social Pediatrics, jointly with the Italian Federation of Pediatricians. *Ital J Pediatr.* 2018;44(1):51. Published 2018 May 8. doi:10.1186/s13052-018-0488-7
- 14. Yeşiltepe Mutlu G, Hatun Ş. Use of Vitamin D in Children and Adults: Frequently Asked Questions. *J Clin Res Pediatr Endocrinol*. 2018;10(4):301 to 306. doi:10.4274/jcrpe.0012

### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care and are solely responsible for the medical advice and treatment of member/enrollees. This clinical policy is not intended to recommend treatment for member/enrollees. Member/enrollees should consult with their treating physician in connection with diagnosis and treatment decisions.



Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, member/enrollees, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, member/enrollees and their representatives agree to be bound by such terms and conditions by providing services to member/enrollees and/or submitting claims for payment for such services.

©2023 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.