

Clinical Policy: Lung Transplantation

Reference Number: LA.CP.MP.57

Date of Last Revision: 04/26

[Coding Implications](#)
[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

This policy describes the medical necessity criteria for the review of lung transplantation requests.

The below criteria are sourced from the International Society for Heart and Lung Transplantation (ISHLT) 2021 Consensus Document for the Selection of Lung Transplant Candidates.¹

The ISHLT consensus document that the below criteria are derived from provides guidelines based on expert synthesis of the current literature with a goal of improving survival and quality of life in transplant candidates. ISHLT recognizes that donor lungs are a limited societal resource, requiring that guidance on candidate selection be based on survival benefit. Given the rigor of the guidelines on which this policy is based, the benefits of receiving a lung transplant in individuals meeting the criteria below outweighs the potential risk of adverse outcomes related to receiving a transplant that is not indicated or not receiving a transplant that is indicated.

Policy/Criteria

- I. It is the policy of Louisiana Healthcare Connections that lung transplantation for members/enrollees with chronic, end-stage lung disease who have failed maximal medical (including pulmonary rehabilitation, as applicable) or surgical therapy is **medically necessary** when all the following criteria are met:
 - A. High (> 50%) risk of death from lung disease within two years if lung transplantation is not performed¹;
 - B. High (> 80%) likelihood of five-year post-transplant survival from a general medical perspective provided there is adequate graft function¹;
 - C. Does not have ANY of the following absolute contraindications¹:
 1. Malignancy with high risk of recurrence or death related to cancer;
 2. Glomerular filtration rate < 40 mL/min/1.73m² unless being considered for multi-organ transplant¹;
 3. Acute renal failure with rising creatinine on dialysis and low likelihood of recovery;
 4. Acute liver failure, or cirrhosis with portal hypertension or synthetic dysfunction unless being considered for multi-organ transplant;
 5. Stroke, acute coronary syndrome, or myocardial infarction (excluding demand ischemia) within 30 days;
 6. Septic shock;
 7. Active extrapulmonary or disseminated infection;
 8. Active *tuberculosis* infection;
 9. HIV infection with detectable viral load unless all of the following are noted:
 - a. CD4 cell count >200 cells/mm³ for at least three months before transplantation²;
 - b. Absence of active AIDS-defining opportunistic infection or malignancy²;
 - c. Member/enrollee is currently on effective antiretroviral therapy (ART)^{2,3,4};

- d. Member/enrollee does not have chronic wasting or severe malnutrition²;
 - 10. Progressive cognitive impairment;
 - 11. Inability to adhere to the regimen necessary to preserve the transplant, even with caregiver support;
 - 12. Other severe, uncontrolled medical condition expected to limit survival after transplant;
 - 13. Active substance use or dependence (including current tobacco use, vaping, marijuana use [unless prescribed by a licensed practitioner], or intravenous drug use) without convincing evidence of risk reduction behaviors (unless urgent transplant timelines are present, in which case a commitment to reducing behaviors is acceptable). Serial blood and urine testing may be used to verify abstinence from substances that are of concern^{1,2};
 - 14. History of nicotine, tobacco, alcohol, or illicit drug use, without documentation noting abstinence from all (including nicotine replacement therapy) for $\geq$ six months prior to transplant²;
- D. Has one of the following disease states (not an all- inclusive list):
- 1. *Adult members/enrollees, age ≥ 18 :*
 - a. Interstitial lung disease and one of the following^{1,*}:
 - i. Absolute decline in forced vital capacity (FVC) $> 10\%$ in the past six months despite appropriate treatment;
 - ii. Absolute decline in diffusing capacity of the lung for carbon monoxide (DLCO) $> 10\%$ in the past six months despite appropriate treatment;
 - iii. Absolute decline in forced vital capacity (FVC) $> 5\%$ with radiographic progression in the past six months despite appropriate treatment;
 - iv. Desaturation to $< 88\%$ on six-minute-walk test (6MWT) or > 50 m decline in 6MWT distance in the past six months;
 - v. Pulmonary hypertension on right heart catheterization or two dimensional echocardiography (in the absence of diastolic dysfunction);
 - vi. Hospitalization because of respiratory decline, pneumothorax, or acute exacerbation;
 - vii. For end-stage or refractory pulmonary alveolar proteinosis (PAP), one of the following:
 - a) Unresponsive to standard therapies, such as whole lung lavage and granulocyte-macrophage colony-stimulating factor (GM-CSF) augmentation;
 - b) Extensive pulmonary fibrosis develops;
 - b. Cystic fibrosis (CF) or other causes of bronchiectasis and any of the following¹:
 - i. $FEV_1 < 25\%$ predicted despite optimal medical management including a trial of elexacaftor/tezacaftor/ivacaftor if eligible;
 - ii. Both of the following:
 - a) Any of the following despite optimal medical management including a trial of elexacaftor/tezacaftor/ivacaftor if eligible:
 - 1) $FEV_1 < 30\%$ predicted;
 - 2) $FEV_1 < 40\%$ predicted and any of the following:
 - i) Six-minute walk distance < 400 meters;
 - ii) $P_aCO_2 > 50$ mmHg;
 - iii) Hypoxemia at rest or with exertion;

- iv) Pulmonary hypertension (PA systolic pressure > 50 mmHg on echocardiogram or evidence of right ventricular dysfunction);
- v) Worsening nutritional status despite supplementation;
- vi) Two exacerbations per year requiring intravenous antibiotics;
- vii) Massive hemoptysis (>240 mL) requiring bronchial artery embolization;
- viii) Pneumothorax;
- 3) FEV₁ <50% predicted and rapidly declining based on pulmonary function testing or progressive symptoms;
- 4) Any exacerbation requiring positive pressure ventilation;
- b) Any of the following¹:
 - 1) Rapid decline in lung function or progressive symptoms (>30% relative decline in FEV₁ over 12 months);
 - 2) Frequent hospitalization, particularly if > 28 days hospitalized in the preceding year;
 - 3) Any exacerbation requiring mechanical ventilation;
 - 4) Chronic respiratory failure with hypoxemia or hypercapnia, particularly for those with increasing oxygen requirements or needing long-term non-invasive ventilation therapy;
 - 5) Pulmonary hypertension (pulmonary arterial systolic pressure >50 mmHg on echocardiogram or evidence of right ventricular dysfunction);
 - 6) Worsening nutritional status particularly with body mass index (BMI) <18 kg/m² despite nutritional interventions;
 - 7) Recurrent massive hemoptysis despite bronchial artery embolization;
 - 8) World Health Organization (WHO) Functional Class IV;
- c. Chronic obstructive pulmonary disease (COPD), and any of the following¹:
 - i. BODE score (includes BMI, degree of airflow obstruction, degree of dyspnea, and exercise capacity) of 7 to 10;
 - ii. FEV₁ (forced expiratory volume in one second) < 20% predicted;
 - iii. History of severe exacerbations;
 - iv. Chronic hypercapnia;
 - v. Moderate to severe pulmonary hypertension;
- d. Pulmonary vascular diseases and any of the following¹:
 - i. European Society of Cardiology/European Respiratory Society (ESC/ERS) high risk or Registry to Evaluate Early and Long-term Pulmonary Arterial Hypertension Disease Management (REVEAL) risk score >10 on appropriate pulmonary arterial hypertension (PAH) therapy, including intravenous (IV) or subcutaneous (SC) prostacyclin analogues;
 - ii. Progressive hypoxemia;
 - iii. Progressive, but not end stage, liver, or kidney dysfunction due to PAH
 - iv. Life-threatening hemoptysis;
 - v. PAH in the European Pediatric Pulmonary Vascular Disease Network (EPPVDN) high risk category and on optimal therapy without improvement¹;
- e. Lymphangioleiomyomatosis (LAM) with evidence of disease progression despite mTOR inhibitor therapy and any of the following¹:
 - i. Severely abnormal lung function (e.g. FEV₁ <30% predicted);
 - ii. Exertional dyspnea (NYHA class III or IV);

- iii. Hypoxemia at rest;
 - iv. Pulmonary hypertension;
 - v. Refractory pneumothorax;
 - f. Primary lung graft failure;
 - g. Acute respiratory distress syndrome (ARDS) with a persistent requirement for mechanical ventilatory support and /or extracorporeal life support (ECLS) without expectation of clinical recovery and with evidence of irreversible lung destruction¹;
2. *Pediatric members/enrollees, age < 18:*
- a. Cystic fibrosis, and any of the following¹:
 - i. $FEV_1 < 30\%$ predicted despite optimal medical management including a trial of elexacaftor/tezacaftor/ivacaftor if eligible¹;
 - ii. Both of the following¹:
 - a) Any of the following despite optimal medical management including a trial of elexacaftor/tezacaftor/ivacaftor if eligible:
 - 1) $FEV_1 < 40\%$ predicted¹;
 - 2) $FEV_1 < 50\%$ predicted and any of the following¹:
 - i) Six-minute walk distance < 400 meters;
 - ii) $P_aCO_2 > 50$ mmHg;
 - iii) Hypoxemia at rest or with exertion;
 - iv) Pulmonary hypertension (PA systolic pressure > 50 mmHg on echocardiogram or evidence of right ventricular dysfunction;
 - v) Worsening nutritional status despite supplementation;
 - vi) Two exacerbations per year requiring intravenous antibiotics;
 - vii) Massive hemoptysis (>240 mL) requiring bronchial artery embolization;
 - viii) Pneumothorax;
 - 3) $FEV_1 < 50\%$ predicted and rapidly declining based on pulmonary function testing or progressive symptoms;
 - 4) Any exacerbation requiring positive pressure ventilation;
 - b) Any of the following¹:
 - 1) Rapid decline in lung function or progressive symptoms (>30% relative decline in FEV_1 over 12 months);
 - 2) Frequent hospitalization, particularly if > 28 days hospitalized in the preceding year;
 - 3) Any exacerbation requiring mechanical ventilation;
 - 4) Chronic respiratory failure with hypoxemia or hypercapnia, particularly for those with increasing oxygen requirements or needing long-term non-invasive ventilation therapy;
 - 5) Pulmonary hypertension (pulmonary arterial systolic pressure >50 mmHg on echocardiogram or evidence of right ventricular dysfunction);
 - 6) Worsening nutritional status particularly with BMI <18 kg/m² despite nutritional interventions;
 - 7) Recurrent massive hemoptysis despite bronchial artery embolization;
 - 8) WHO Functional Class IV;
 - b. Pulmonary vascular disease and any of the following¹:
 - i. ESC/ERS high risk or Registry to Evaluate Early and Long-term Pulmonary Arterial Hypertension Disease Management (REVEAL) risk

- score >10 on appropriate PAH therapy, including IV or SC prostacyclin analogues;
- ii. Progressive hypoxemia;
- iii. Progressive, but not end stage, liver, or kidney dysfunction due to PAH;
- iv. Life-threatening hemoptysis;
- c. Interstitial lung disease and one of the following, any of the following¹:
 - i. Absolute decline in FVC > 10% in the past six months despite appropriate treatment;
 - ii. Absolute decline in DLCO > 10% in the past six months despite appropriate treatment;
 - iii. Absolute decline in FVC > 5% with radiographic progression in the past six months despite appropriate treatment;
 - iv. Desaturation to < 88% on 6MWT or > 50 m decline in 6MWT distance in the past six months;
 - v. Pulmonary hypertension on right heart catheterization or two dimensional echocardiography (in the absence of diastolic dysfunction);
 - vi. Hospitalization because of respiratory decline, pneumothorax, or acute exacerbation;
 - vii. For end-stage or refractory pulmonary alveolar proteinosis (PAP), one of the following:
 - a) Unresponsive to standard therapies, such as whole lung lavage and granulocyte-macrophage colony-stimulating factor (GM-CSF) augmentation;
 - b) Extensive pulmonary fibrosis develops;
- d. COPD, and any of the following¹:
 - i. BODE score (includes BMI, degree of airflow obstruction, degree of dyspnea, and exercise capacity) of 7 to 10;
 - ii. FEV₁ < 20% predicted;
 - iii. History of severe exacerbations;
 - iv. Chronic hypercapnia;
 - v. Moderate to severe pulmonary hypertension;
- e. Primary lung graft failure¹;
- f. LAM with evidence of disease progression despite mTOR inhibitor therapy and any of the following¹:
 - i. Several abnormal lung function (e.g. FEV₁ <30% predicted);
 - ii. Exertional dyspnea (NYHA class III or IV);
 - iii. Hypoxemia at rest;
 - iv. Pulmonary hypertension;
 - v. Refractory pneumothorax;
- g. ARDS with a persistent requirement for mechanical ventilatory support and /or ECLS without expectation of clinical recovery and with evidence of irreversible lung destruction¹;
- h. Alveolar capillary dysplasia¹;
- i. Pulmonary vein stenosis refractory to intervention¹;
- j. Pulmonary veno-occlusive disease.¹

**Note:* FVC may be a less reliable parameter for those with concomitant emphysema.¹

Background

Lung transplantation is an accepted therapy for the management of a range of severe lung disorders. Single, double, and lobar-lung transplants have all been successful for carefully selected patients with end-stage pulmonary disease. The most common disease processes for which lung transplants are performed include chronic obstructive pulmonary disease (COPD), idiopathic pulmonary fibrosis, cystic fibrosis, pulmonary arterial hypertension, and sarcoidosis.⁵

COPD is one of the most common lung diseases and is the most common indication for lung transplantation in adults. Chronic bronchitis and emphysema are the two main forms of COPD, both most commonly caused by smoking. Non-smokers with an alpha-1 antitrypsin deficiency can also develop emphysema. These conditions are the most common indications for single lung transplants. Cystic fibrosis, emphysema, and alpha-1 antitrypsin deficiency are the most common indications for double lung transplant, or sequential replacement of both lungs.

The most common indications for pediatric lung transplants include pulmonary vascular disease, bronchiolitis obliterans, bronchopulmonary dysplasia, graft failure due to viral pneumonitis, and cystic fibrosis.

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2025, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only and may not support medical necessity and may not support medical necessity. Inclusion or exclusion of any codes does not guarantee coverage.

Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

NOTE: Coverage is subject to each requested code's inclusion on the corresponding LDH fee schedule. Non-covered codes are denoted (*) and are reviewed for Medical Necessity for members under 21 years of age on a per case basis.

CPT® Codes	Description
32850	Donor pneumonectomy(s) (including cold preservation), from cadaver donor
32851	Lung transplant, single; without cardiopulmonary bypass
32852	Lung transplant, single; with cardiopulmonary bypass
32853	Lung transplant, double (bilateral sequential or en bloc); without cardiopulmonary bypass
32854	Lung transplant, double (bilateral sequential or en bloc); with cardiopulmonary bypass
32855*	Backbench standard preparation of cadaver donor lung allograft prior to transplantation, including dissection of allograft from surrounding soft tissues to prepare pulmonary venous/atrial cuff, pulmonary artery, and bronchus; unilateral

CPT® Codes	Description
32856*	Backbench standard preparation of cadaver donor lung allograft prior to transplantation, including dissection of allograft from surrounding soft tissues to prepare pulmonary venous/atrial cuff, pulmonary artery, and bronchus; bilateral

HCPCS Codes	Description
S2060*	Lobar lung transplantation
S2152*	Solid organ(s), complete or segmental, single organ or combination of organs; deceased or living donor (s), procurement, transplantation, and related complications; including: drugs; supplies; hospitalization with outpatient follow-up; medical/surgical, diagnostic, emergency, and rehabilitative services, and the number of days of pre- and post-transplant care in the global definition

Reviews, Revisions, and Approvals	Revision Date	Approval Date	Effective Date
Converted corporate to local policy.	08/15/20		
References reviewed and updated. Replaced “members” with “members/enrollees” in all instances. Replaced contraindications of “severely limited functional status with poor rehabilitation potential” and those regarding past or current nonadherence to medical therapy, and psychological condition associated with the inability to comply with medical therapy with “Inability to adhere to the regimen necessary to preserve the transplant, even with caregiver support.” Changed “review date” in header to “Date of Last Revision” and “Date” in the revision log header to “Revision Date.” Added “and may not support medical necessity” in coding implications. Annual review. References reviewed and updated. Reviewed by specialist.	2/22		
Annual review. Added “or surgical therapy” to I and noted that maximal medical therapy includes pulmonary rehab when applicable. Updated the following based on ISHLT 2021 guidelines; removed criteria “High (> 80%) likelihood of surviving at least 90 days after lung transplantation.”, updated I.C., I.D.1.a, I.D.1.b., I.D.1.c., I.D.1.d., I.D.1.f., I.D.2.a, I.D.2.b. Clarified nicotine and tobacco abstinence contraindication. Added CPT codes 32850, 32855, and 32856. References reviewed, updated, and reformatted. Reviewed by specialist. Added “and may not support medical necessity” to Coding Implications section	5/22	8/13/22	
Annual review. Criteria I.C.14. updated to exclude marijuana use when prescribed by a licensed practitioner and include required commitment to reducing substance use behaviors if urgent transplant timelines are present. Added pediatric indication for end-stage emphysema due to alpha-1 trypsin deficiency. ICD-10 codes removed. References	4/23	7/21/23	

Reviews, Revisions, and Approvals	Revision Date	Approval Date	Effective Date
reviewed and updated. Reviewed by external specialist.			
Revised adult and pediatric criteria to align with ISHLT 2021 consensus document. References reviewed and updated. Note for non-covered codes added.	09/23	11/27/23	12/27/23
Annual review. Updated I.C.2. from GFR < 40 mL/min/1.73m ² to GFR < 30 mL/min/1.73m ² . Expanded I.C.9. with qualifying criteria for members who are HIV positive. Updated I.D.2.a.1. from FEV ₁ <25% to FEV ₁ <30%. Background updated with no impact to criteria. References reviewed and updated.	05/24	7/16/24	8/16/24
Annual review. Updated glomerular filtration rate from < 30 to < 40 mL/min/1.73m ² in Criteria I.C.2. Updated Criteria I.C.9.a. to include at least three months prior to transplantation. Removed additional information regarding heart transplant waiting list in Criteria I.C.9.b. Minor grammatical update in Criteria I.C.9.c. Added Criteria I.C.9.d. regarding chronic wasting or severe malnutrition. Expanded Criteria I.C.13. regarding active substance use or dependence and added Criteria I.C.14. regarding documentation of abstinence from substance use. Minor grammatical changes to Criteria I.D.1.b.ii.b)5), Criteria I.D.1.c.i., Criteria I.D.2., Criteria I.D.2.a.ii.b)5), and Criteria I.D.2.d.i. with no clinical significance. Added Criteria I.D.2.h., Criteria I.D.2.i, and Criteria I.D.2.j. regarding alveolar capillary dysplasia, pulmonary vein stenosis refractory to intervention, and pulmonary veno-occlusive disease. Background updated with no impact to criteria. References reviewed and updated. Reviewed by internal specialist and external specialist.	3/25	5/20/25	6/19/25
Annual review. Updated adult and pediatric interstitial lung disease criteria to include end-stage or refractory pulmonary alveolar proteinosis as criteria I.D.1.c.vii.a)-b) and I.D.2.c.vii.a)-b) respectively. Reviewed codes and descriptions. References reviewed and updated.	04/26	7/7/26	8/7/26

References

1. Leard LE, Holm AM, Valapour M, et al. Consensus document for the selection of lung transplant candidates: An update from the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant*. 2021;40(11):1349 to 1379. doi:10.1016/j.healun.2021.07.005
2. Peled Y, Ducharme A, Kittleson M, et al. International Society for Heart and Lung Transplantation Guidelines for the Evaluation and Care of Cardiac Transplant Candidates-2024. *J Heart Lung Transplant*. 2024;43(10):1529-1628.e54. doi:10.1016/j.healun.2024.05.010
3. Agüero F, Castel MA, Cocchi S, et al. An Update on Heart Transplantation in Human Immunodeficiency Virus-Infected Patients. *Am J Transplant*. 2016;16(1):21-28. doi:10.1111/ajt.13496
4. Harbell J, Terrault NA, Stock P. Solid organ transplants in HIV-infected patients. *Curr HIV/AIDS Rep*. 2013;10(3):217-225. doi:10.1007/s11904-013-0170-z
5. Hachem RR. Lung transplantation: an overview. UpToDate. www.uptodate.com.

Published May 23, 2024. Accessed December 15, 2025.

6. Yusen RD, Edwards LB, Kucheryavaya AY, et al. The Registry of the International Society for Heart and Lung Transplantation: Thirty-second Official Adult Lung and Heart-Lung Transplantation Report--2015; Focus Theme: Early Graft Failure. *J Heart Lung Transplant*. 2015;34(10):1264-1277. doi:10.1016/j.healun.2015.08.014
7. Faro A, Mallory GB, Visner GA, et al. American Society of Transplantation executive summary on pediatric lung transplantation. *Am J Transplant*. 2007;7(2):285 to 292. doi:10.1111/j.1600-6143.2006.01612.x
8. Hachem RR. Lung transplantation: disease-based choice of procedure. UpToDate. www.uptodate.com. Published February 14, 2025. Accessed December 15, 2025.
9. Hachem RR. Lung transplantation: general guidelines for recipient selection. UpToDate. www.uptodate.com. Published October 30, 2025. Accessed December 15, 2025.
10. Hall DJ, Belli EV, Gregg JA, et al. Two Decades of Lung Retransplantation: A Single-Center Experience. *Ann Thorac Surg*. 2017;103(4):1076-1083. doi:10.1016/j.athoracsur.2016.09.107
11. Kirkby S, Hayes D Jr. Pediatric lung transplantation: indications and outcomes. *J Thorac Dis*. 2014;6(8):1024-1031. doi:10.3978/j.issn.2072-1439.2014.04.27
12. Kotloff RM, Thabut G. Lung transplantation. *Am J Respir Crit Care Med*. 2011;184(2):159-171. doi:10.1164/rccm.201101-0134CI
13. Meyer KC. Recent advances in lung transplantation. *F1000Res*. 2018;7:F1000 Faculty Rev-1684. Published 2018 Oct 23. doi:10.12688/f1000research.15393.1
14. Whitson, BA. Lung transplantation. Medscape. <https://emedicine.medscape.com/article/429499-overview>. Published August 19, 2019. Updated March 14, 2025. Accessed December 12, 2025.
15. National Institute for Health and Care Excellence. Living-donor lung transplantation for end-stage lung disease. Interventional procedures guidance [IPG170]. <https://www.nice.org.uk/guidance/ipg170>. Published May 24, 2006. Accessed December 12, 2025.
16. Organ Procurement and Transplantation Network. Policies. <https://optn.transplant.hrsa.gov/policies-bylaws/policies/>. Updated December 11, 2024. Accessed December 12, 2025.
17. Simon, RH. Cystic fibrosis: management of advanced lung disease. UpToDate. www.uptodate.com. Published December 10, 2024. Accessed December 15, 2025.
18. MedlinePlus. Chronic obstructive pulmonary disease (COPD). <https://medlineplus.gov/ency/article/000091.htm>. Published May 03, 2023. Accessed December 12, 2025.
19. Stone HM, Edgar RG, Thompson RD, Stockley RA. Lung Transplantation in Alpha-1-Antitrypsin Deficiency. *COPD*. 2016;13(2):146-152. doi:10.3109/15412555.2015.1048850
20. Stoller JK. Clinical manifestations, diagnosis, and natural history of alpha-1 antitrypsin deficiency. UpToDate. www.uptodate.com. Published October 01, 2024. Accessed December 15, 2025.
21. Stoller JK. Treatment of emphysema from alpha-1 antitrypsin deficiency. UpToDate. www.uptodate.com. Published April 25, 2025. Accessed December 15, 2025.
22. Roest S, Hesselink DA, Klimczak-Tomaniak D, et al. Incidence of end-stage renal disease after heart transplantation and effect of its treatment on survival. *ESC Heart Fail*.

- 2020;7(2):533-541. doi:10.1002/ehf2.12585
23. Biswas Roy S, Panchanathan R, Walia R, et al. Lung Retransplantation for Chronic Rejection: A Single-Center Experience. *Ann Thorac Surg*. 2018;105(1):221-227. doi:10.1016/j.athoracsur.2017.07.025
24. Christie JD, Edwards LB, Kucheryavaya AY, et al. The Registry of the International Society for Heart and Lung Transplantation: Twenty-eighth Adult Lung and Heart-Lung Transplant Report--2011. *J Heart Lung Transplant*. 2011;30(10):1104-1122. doi:10.1016/j.healun.2011.08.004
25. Rabe KF, Watz H. Chronic obstructive pulmonary disease. *Lancet*. 2017;389(10082):1931-1940. doi:10.1016/S0140-6736(17)31222-9
26. Lee JS, Montesi S. Overview of the management of adults with interstitial lung disease. UpToDate. www.uptodate.com. Published December 16, 2025. Accessed December 19, 2025.
27. Hayes D Jr, Avdimiretz N, Brugha R, et al. International Society for Heart and Lung Transplantation Consensus Statement on the Referral and Selection of Pediatric Lung Transplant Candidates. *J Heart Lung Transplant*. 2025;44(12):e133-e176. doi:10.1016/j.healun.2025.08.005

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise

professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members/enrollees. This clinical policy is not intended to recommend treatment for members/enrollees. Members/enrollees should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members/enrollees and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members/enrollees and their representatives agree to be bound by such terms and conditions by providing services to members/enrollees and/or submitting claims for payment for such services.

©2026 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.