

## Clinical Policy: Testing for Select Genitourinary Conditions

Reference Number: LA.CP.MP.97

Date of Last Revision: 1/2022

Coding Implications

Revision Log

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### Description

Various diagnostic methods are available to identify the etiology of the signs and symptoms of vaginitis. The purpose of this policy is to define medical necessity criteria for the diagnostic evaluation of vaginitis (*excluding Trichomonas vaginalis, vaginal pH testing, and microscopic examination with saline and KOH*) in members/enrollees  $\geq 13$  years of age. This policy also defines unnecessary amplified DNA- (deoxyribonucleic acid) probe testing for genitourinary conditions.

### Policy/Criteria

- I. It is the policy of Louisiana Healthcare Connections that the following diagnostic tests for symptomatic women for the evaluation of vaginitis are medically necessary for members/enrollees age  $\geq 13$ :
  - A. Vaginal pH, microscopic examination with 0.9% saline and 10% KOH, and KOH “whiff test (i.e., amine odor test);
  - B. Assay for sialidase activity;
  - C. Direct DNA probe tests to detect the presence of *Candida and Gardnerella vaginalis*.
  
- II. It is the policy of Louisiana Healthcare Connections that screening of asymptomatic pregnant women for bacterial vaginosis (BV) to reduce the incidence of pre-term birth or other complications of pregnancy is not medically necessary as there is no evidence that treatment of BV in asymptomatic pregnant women reduces these complications.<sup>9</sup>
  
- III. It is the policy of Louisiana Healthcare Connections that unspecified amplified DNA-probe testing for genitourinary conditions for asymptomatic women during routine exams, contraceptive management care, or pregnancy care is considered not medically necessary for members/enrollees  $\geq 13$  year of age as it has not been shown to improve clinical outcomes over direct DNA-probe testing.
  
- IV. It is the policy of Louisiana Healthcare Connections that unspecified amplified DNA-probe testing for the diagnostic evaluation of symptomatic women for the following genitourinary conditions is considered not medically necessary for members/enrollees  $\geq 13$  of age as it has not been shown to improve clinical outcomes over direct DNA-probe testing:
  - A. Acute vaginitis or vulvitis ( $\leq 4$  episodes per year);
  - B. Gynecologic and obstetric conditions triggered by etiologies other than complicated vaginitis inducing mechanisms as listed in Table 5, including:
    1. Urinary tract infections;
    2. Pelvic inflammatory disease;
    3. Inflammatory disorders of the vagina, vulva, and perineum;
    4. Irregular menstruation or abnormal uterine and vaginal bleeding;
    5. Dysmenorrhea;
    6. Complications with pregnancy, including all of the following:

- a. Pre-term labor;
- b. Ectopic pregnancy;
- c. High risk pregnancy.

### **Background**

The 3 diseases most frequently associated with vaginitis are BV, (caused by replacement of the vaginal flora by an overgrowth of anaerobic bacteria, including *Prevotella* sp., *Mobiluncus* sp., *Gardnerella vaginalis*, *Ureaplasma*, *Mycoplasma*, and numerous fastidious or uncultivated anaerobes), trichomoniasis (caused by *Trichomonas vaginalis*), and candidiasis (usually caused by *Candida albicans*).<sup>3</sup>

Various diagnostic methods are available to identify the etiology of the signs and symptoms of vaginitis. The cause of vaginal symptoms might be determined by testing of pH, for the presence of amines by the use of a potassium hydroxide (KOH) test, and microscopic examination of fresh samples of the discharge. The pH of the vaginal secretions can be determined by narrow-range pH paper; an elevated pH (*i.e.*, >4.5) is common with BV or trichomonas. Because pH testing is not highly specific, the discharge should be further examined microscopically with both a saline and KOH solution.<sup>3</sup>

The saline-solution specimen might yield motile *T. vaginalis* or clue cells (*i.e.*, epithelial cells with borders obscured by small bacteria), which are characteristic of BV, whereas the presence of white blood cells without evidence of trichomonads or yeast in this solution is suggestive of cervicitis. The KOH specimen typically is used to identify the yeast or pseudohyphae of *Candida* species. However, the absence of trichomonads or pseudohyphae in KOH samples does not rule out these infections, because the sensitivity of microscopy is approximately 40-75%.<sup>4</sup>

In settings where pH paper, KOH, and microscopy are not available or are inconclusive, alternative, commercially available, point-of-care tests, such as commercially available direct DNA-probe tests, or clinical laboratory testing, can be used to diagnose vaginitis. The presence of objective signs of vulvar inflammation in the absence of vaginal pathogens after laboratory testing, along with a minimal amount of discharge, suggests the possibility of mechanical, chemical, allergic, or other noninfectious irritation of the vulva.<sup>5</sup>

### *Bacterial Vaginosis*

BV is a polymicrobial clinical syndrome resulting from replacement of the normal hydrogen peroxide-producing *Lactobacillus* species in the vagina with high concentrations of anaerobic bacteria (*e.g.*, *Prevotella* sp. and *Mobiluncus* sp.), *G. vaginalis*, *Ureaplasma*, *Mycoplasma*, and numerous fastidious or uncultivated anaerobes. BV is the most prevalent cause of vaginal discharge or malodor; however, in a nationally representative survey, most women with BV were asymptomatic.<sup>3</sup>

BV can be diagnosed by the use of clinical criteria (*i.e.*, Amsel's Diagnostic Criteria) or Gram stain (considered the gold standard laboratory method for diagnosing BV). If a Gram stain is not available, clinical criteria can be used and require 3 of the following symptoms or signs:

- Homogeneous, thin, white discharge that smoothly coats the vaginal walls;
- Presence of clue cells on microscopic examination;

- pH of vaginal fluid >4.5;
- A fishy odor of vaginal discharge before or after addition of 10% KOH (*i.e.*, the whiff test).<sup>8</sup>

Detection of 3 of these criteria has been correlated with results by Gram stain. Other tests, including a DNA probe-based test for high concentrations of *G. vaginalis* (Affirm VP III, Becton Dickinson, Sparks, Maryland), and the OSOM BVBlue test have acceptable performance characteristics compared with Gram stain.<sup>3</sup> The BVBlue test is a colorimetric test for enzymes (sialidase) produced by BV organisms. Culture of *G. vaginalis* is not recommended as a diagnostic tool, because 50% of women in the general population will culture positively for *G. vaginalis*. Use of the proline-aminopeptidase test card (Pip Activity TestCard) is no longer recommended because of low sensitivity and specificity.<sup>8</sup>

#### *Vulvovaginal Candidiasis*

Vulvovaginal candidiasis (VVC) usually is caused by *C. albicans*, but occasionally is caused by other *Candida* species or yeasts. Typical symptoms of VVC include pruritus, vaginal soreness, dyspareunia, external dysuria, and abnormal vaginal discharge. None of these symptoms is specific for VVC. An estimated 75% of women will have at least 1 episode of VVC, and 40%–45% will have 2 or more episodes within their lifetime. On the basis of clinical presentation, microbiology, host factors, and response to therapy, VVC can be classified as either uncomplicated or complicated.<sup>3</sup>

A diagnosis of *Candida* vaginitis is suggested clinically by the presence of external dysuria and vulvar pruritus, pain, swelling, and redness. Signs include vulvar edema, fissures, excoriations, or thick, curdy vaginal discharge. The diagnosis can be made in a woman who has signs and symptoms of vaginitis when either 1) a wet preparation (saline, 10% KOH) or Gram stain of vaginal discharge demonstrates yeasts, hyphae, or pseudohyphae or 2) a culture or other test yields a yeast species. *Candida* vaginitis is associated with a normal vaginal pH (<4.5), and therefore, pH testing is not a useful diagnostic tool. Use of 10% KOH in wet preparations improves the visualization of yeast and mycelia by disrupting cellular material that might obscure the yeast or pseudohyphae. Examination of a wet mount with KOH preparation should be performed for all women with symptoms or signs of VVC, and women with a positive result should receive treatment. For women with negative wet mounts who are symptomatic, vaginal cultures for *Candida* should be considered. If the wet mount is negative and *Candida* cultures cannot be done, empiric treatment can be considered for symptomatic women with any sign of VVC on examination. Identifying *Candida* by culture in the absence of symptoms or signs is not an indication for treatment because approximately 10%–20% of women harbor *Candida* species and other yeasts in the vagina. VVC can occur concomitantly with STIs. Most healthy women with uncomplicated VVC have no identifiable precipitating factors.<sup>3</sup>

Complicated or recurrent vulvovaginal candidiasis (RVVC) is usually defined as 4 or more episodes of symptomatic VVC in 1 year, and affects a small percentage of women (<5%). The pathogenesis of RVVC is poorly understood, and most women with RVVC have no apparent predisposing or underlying conditions. Vaginal cultures should be obtained from patients with RVVC to confirm the clinical diagnosis and to identify unusual species (including nonalbicans species), particularly *Candida glabrata*. Although *C. glabrata* and other nonalbicans *Candida*

species are observed in 10%-20% of patients with RVVC, *C. glabrata* does not form pseudohyphae or hyphae and is not easily recognized on microscopy.<sup>3</sup>

VVC occurs more frequently and has greater persistence, but not greater severity, in HIV- (human immunodeficiency virus) infected women with very low CD4 counts and high viral load. However, this population is likely to manifest other acquired immune deficiency syndrome – related sentinel conditions. HIV testing of women only for the indication of recurrent vulvovaginal candidiasis is not justified, given that this condition is common in women without HIV.<sup>8</sup>

DNA-probe tests have been developed to directly detect the presence of *Candida*, *Trichomonas* and *G. vaginalis*. Since *G. vaginalis* is a normal part of the vaginal flora, the DNA probe test is designed to be relatively insensitive, detecting only pathogenic levels of *G. vaginalis*. DNA probes amplified by PCR testing can also detect these pathogens. In PCR tests, the sample is treated with enzymes that amplify specific regions of the DNA. After amplification, the number of DNA fragments is quantified. PCR testing has proven to be the most accurate diagnostic method in recent studies; however PCR testing has not been shown to improve clinical outcomes over direct DNA-probe testing.

#### *Pediatric Patients*

Girls less than 13 years of age tend to have a different etiology for vaginitis than do older girls, due to the lack of estrogenization of the vagina, and the consequential alkalinity and vaginal atrophy.<sup>5</sup> Common causes of vulvovaginal symptoms may include respiratory organisms such as group A streptococci and *Hemophilus influenzae*, as well as enteric and sexually transmitted pathogens. Pinworms or foreign bodies may also lead to vaginitis in this population..<sup>5 6</sup>

#### *Centers for Disease Control and Prevention*

Screening for *T. vaginalis* in women can be considered in those at high risk for infection (i.e., women who have new or multiple partners, have a history of STIs, exchange sex for payment, and use injection drugs).<sup>3</sup>

Recommends the gram stain as the gold standard for diagnosis of bacterial vaginosis, and recommend use of Amsel's criteria if a gram stain is not available.<sup>3</sup>

#### *U.S. Preventive Services Task Force*

This organization does not recommend screening for bacterial vaginosis in pregnant women at low-risk for pre-term delivery.<sup>9</sup>

#### **Coding Implications**

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2019, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only and may not support medical necessity. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources

of professional coding guidance prior to the submission of claims for reimbursement of covered services.

Table 1. CPT codes considered medically necessary when billed with an ICD-10-CM code in Table 2

<b>CPT®*</b> <b>Codes</b>	<b>Description</b>
82120	Amines, vaginal fluid, qualitative
87480	Infectious agent detection by nucleic acid (DNA or RNA); Candida species, direct probe technique
87510	Infectious agent detection by nucleic acid (DNA or RNA); Gardnerella vaginalis, direct probe technique
87905	Infectious agent enzymatic activity other than virus (eg, sialidase activity in vaginal fluid)

Table 2. ICD-10-CM diagnosis codes that support medical necessity per this policy

<b>ICD-10-CM</b> <b>Code</b>	<b>Description</b>
B37.3	Candidiasis of vulva and vagina
L29.2, L29.3	Pruritus of genitals
N76.0 – N76.3	Vaginitis and vulvitis
N77.1	Vaginitis, vulvitis, and vulvovaginitis in diseases classified elsewhere
N89.8	Other specific noninflammatory disorders of vagina
O23.511– O23.93	Infection of genitourinary tract in pregnancy
Z72.51 – Z72.53	High risk sexual behavior [exchange of sex for payment, new or multiple partners]
Z86.19	Personal history of other infectious and parasitic diseases [history of STDs]

Table 3. CPT codes considered not medically necessary unless an exception is noted in this policy.

CPT Codes	Description
81513	Infectious disease, bacterial vaginosis, quantitative real-time amplification of RNA markers for <i>Atopobium vaginae</i> , <i>Gardnerella vaginalis</i> , and <i>Lactobacillus</i> species, utilizing vaginal-fluid specimens, algorithm reported as a positive or negative result for bacterial vaginosis
81514	Infectious disease, bacterial vaginosis and vaginitis, quantitative real-time amplification of DNA markers for <i>Gardnerella vaginalis</i> , <i>Atopobium vaginae</i> , <i>Megasphaera</i> type 1, Bacterial Vaginosis Associated Bacteria-2 (BVAB-2), and <i>Lactobacillus</i> species ( <i>L. crispatus</i> and <i>L. jensenii</i> ), utilizing vaginal-fluid specimens, algorithm reported as a positive or negative for high likelihood of bacterial vaginosis, includes separate detection of <i>Trichomonas vaginalis</i> and/or <i>Candida</i> species ( <i>C. albicans</i> , <i>C. tropicalis</i> , <i>C. parapsilosis</i> , <i>C. dubliniensis</i> ), <i>Candida glabrata</i> , <i>Candida krusei</i> , when reported
87481	Infectious agent detection by nucleic acid (DNA or RNA); <i>Candida</i> species, amplified probe technique
87511	Infectious agent detection by nucleic acid (DNA or RNA); <i>Gardnerella vaginalis</i> , amplified probe technique

Table 4. CPT codes considered not medically necessary when billed with an ICD-10-CM code listed in Table 5 below.

CPT Codes	Description
87798	Infectious agent detection by nucleic acid (DNA or RNA), not otherwise specified; amplified probe technique, each organism

Table 5. ICD-10-CM diagnosis codes considered not medically necessary when billed with CPT code 87798 per this policy.

ICD-10-CM Code	Description
N39.0	Urinary tract infection, site not specified
N72	Inflammatory disease of cervix uteri
N76.0	Acute vaginitis
N76.2	Acute vulvitis
N89.9	Noninflammatory disorder of vagina, unspecified
N90.89	Other specified noninflammatory disorders of vulva and perineum
N90.9	Noninflammatory disorder of vulva and perineum, unspecified
N91.0 – N91.5	Absent, scanty and rare menstruation
N92.0	Excessive, frequent menstruation with regular cycle
N93.0	Postcoital and contact bleeding
N93.8	Other specified abnormal uterine and vaginal bleeding
N93.9	Abnormal uterine and vaginal bleeding, unspecified
N94.3	Premenstrual tension syndrome
N94.4 – N94.6	Dysmenorrhea

ICD-10-CM Code	Description
N94.89	Other specified conditions associated with female genital organs and menstrual cycle
N94.9	Unspecified condition associated with female genital organs and menstrual cycle
O09.00-O09.03	Supervision of pregnancy with history of infertility
O09.10-O09.13	Supervision of pregnancy with history of ectopic pregnancy
O09.A0-O09.A3	Supervision of pregnancy with history of molar pregnancy
O09.211-O09.219	Supervision of pregnancy with history of pre-term labor
O09.291-O09.299	Supervision of pregnancy with other poor reproductive or obstetric history
O09.30-O09.33	Supervision of pregnancy with insufficient antenatal care
O09.40-O09.43	Supervision of pregnancy with grand multiparity
O09.511-O09.519	Supervision of elderly primigravida
O09.521- O09.529	Supervision of elderly multigravida
O09.611-O09.619	Supervision of young primigravida
O09.621-O09.629	Supervision of young multigravida
O09.70-O09.73	Supervision of high risk pregnancy due to social problems
O09.811-O09.819	Supervision of pregnancy resulting from assisted reproductive technology
O09.821-O09.829	Supervision of pregnancy with history of in utero procedure during previous pregnancy
O09.891-O09.899	Supervision of other high risk pregnancies
O09.90-O09.93	Supervision of high risk pregnancy, unspecified
Z00.00	Encounter for general adult medical examination without abnormal findings
Z00.8	Encounter for other general examination
Z01.419	Encounter for gynecological examination (general) (routine) without abnormal findings
Z11.3	Encounter for screening for infections with a predominantly sexual mode of transmission
Z11.51	Encounter for screening for human papillomavirus (HPV)
Z22.330	Carrier of Group B streptococcus
Z23	Encounter for immunization
Z30.011 – Z30.019	Encounter for initial prescription of contraceptives
Z30.02	Counseling and instruction in natural family planning to avoid pregnancy
Z30.09	Encounter for other general counseling and advice on contraception
Z30.40 – Z30.9	Encounter for surveillance of contraceptives
Z32.00	Encounter for pregnancy test, result unknown
Z33.1	Pregnant state, incidental
Z34.00 – Z34.03	Encounter for supervision of normal first pregnancy
Z34.80 – Z34.83	Encounter for supervision of other normal pregnancy
Z34.90 – Z34.93	Encounter for supervision of normal pregnancy, unspecified
Z36.0-Z36.5	Encounter for antenatal screening of mother

ICD-10-CM Code	Description
Z36.81-Z36.9	Encounter for other antenatal screening
Z38.00 – Z38.01	Single liveborn infant, born in hospital
Z38.30 – Z38.31	Twin liveborn infant, born in hospital
Z38.61 – Z38.69	Other multiple liveborn infant, born in hospital
Z39.0 – Z39.2	Encounter for maternal postpartum care and examination
Z3A.00 – Z3A.49	Weeks of gestation
Z97.5	Presence of (intrauterine) contraceptive device

Reviews, Revisions, and Approvals	Revision Date	Approval Date
Converted corporate to local policy.	08/15/2020	
Noted in the description that the policy does not apply to the diagnosis of Trichomonas vaginalis, vaginal pH testing, and wet mount microscope tests, and updated background accordingly. Changed “review date” in the header to “date of last revision” and “date” in the revision log header to “revision date.” References reviewed, reformatted and updated. Removed 83986 and 87210 from the coding table requiring symptom diagnosis codes, as they could be used for testing for conditions other than vaginitis. Removed the following codes from table 2: A59.01, F11.10 - F11.19, F11.20 – F11.29, F14.10 – F14.19, F14.20 – F14.29, F15.10 – F15.19, F15.20 – F15.29, F18.10 – F18.19, F18.20 – F18.29, F19.10 – F19.19, F19.20 – F19.29, Z11.2, Z11.8, Z13.89. Specialist review.	1/2022	

**References**

1. Current Procedural Terminology (CPT®), 2021
2. ICD-10-CM Official Code Set, 2021
3. Workowski KA, Bolan GA; Centers for Disease Control and Prevention Sexually transmitted diseases treatment guidelines, 2015 [published correction appears in MMWR Recomm Rep. 2015 Aug 28;64(33):924]. MMWR Recomm Rep. 2015;64(RR-03):1-137.
4. Ferris DG, Hendrich J, Payne PM, et al. Office laboratory diagnosis of vaginitis. Clinician-performed tests compared with a rapid nucleic acid hybridization test. *J Fam Pract.* 1995;41(6):575-581.
5. Vaginitis in Nonpregnant Patients: ACOG Practice Bulletin, Number 215. *Obstet Gynecol.* 2020;135(1):e1-e17. doi:10.1097/AOG.0000000000003604
6. Kaufman CA. Overview of Candida infections. UpToDate Website. [www.uptodate.com](http://www.uptodate.com). Published May 26, 2021. Accessed July 2, 2021.
7. Kinny RG, Spach DH. Vaginitis. National Sexually Transmitted Disease Curriculum. February 2017. Updated Oct 26, 2018. Available at: <http://www.std.uw.edu/go/syndrome-based/vaginal-discharge/core-concept/all#summary-points>
8. Sobel JD. Approach to females with symptoms of vaginitis. UpToDate Website. [www.uptodate.com](http://www.uptodate.com). Published May 25, 2021. Accessed July 2, 2021.



9. U.S. Preventive Services Task Force, Owens DK, Davidson KW, et al. Bacterial Vaginosis in Pregnant Persons to Prevent Preterm Delivery: US. Preventive Services Task Force Recommendation Statement. *JAMA*. 2020;323(13):1286-1292. doi: 10.1001/jama.2020.2684.
10. Vaginitis Panels by Direct DNA Probe. Diagnostic Laboratory Services. October 2012.
11. Van Der Pol B. Clinical and Laboratory Testing for *Trichomonas vaginalis* Infection. *J Clin Microbiol*.2016;54(1):7-12. doi:10.1128/JCM.02025-15
12. Sobol JD, Mitchell C. Trichomoniasis. UpToDate website. [www.uptodate.com](http://www.uptodate.com). Published April 20, 2021. Accessed July 02, 2021.
13. Schwebke JR, Taylor SN, Ackerman R, et al. Clinical Validation of the Aptima Bacterial Vaginosis and Aptima *Candida/Trichomonas* Vaginitis Assays: Results from a Prospective Multicenter Clinical Study. *J Clin Microbiol*. 2020 Jan 28. doi: 10.1128/JCM.01643-19.
14. Laufer MR, Emans SJ. Overview of vulvovaginal complaints in the prepubertal child. UpToDate website. [www.uptodate.com](http://www.uptodate.com). Published April 16, 2020. Accessed July 6, 2021.

### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise

## CLINICAL POLICY

### Testing for Select GU Conditions



professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members/enrollees. This clinical policy is not intended to recommend treatment for members/enrollees. Members/enrollees should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members/enrollees and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members/enrollees and their representatives agree to be bound by such terms and conditions by providing services to members/enrollees and/or submitting claims for payment for such services.

©2020 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.