

Clinical Policy: Ziv-Aflibercept (Zaltrap)

Reference Number: LA.PHAR.325

Effective Date: 11.04.23

Last Review Date: 09.21.26

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

****Please note: This policy is for medical benefit****

Description

Ziv-aflibercept (Zaltrap®) is a vascular endothelial growth factor (VEGF) inhibitor.

FDA Approved Indication(s)

Zaltrap, in combination with 5-fluorouracil, leucovorin, irinotecan (FOLFIRI), is indicated for patients with metastatic colorectal cancer (CRC) that is resistant to or has progressed following an oxaliplatin-containing regimen.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections that Zaltrap is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Colorectal Cancer (must meet all):

1. Diagnosis of advanced, unresectable, or metastatic CRC;
2. Prescribed by or in consultation with an oncologist;
3. Age $\geq$ 18 years;
4. Previous treatment with one of the following (a, b, or c):
 - a. An oxaliplatin-containing regimen (e.g., FOLFOX, CapeOX);
 - b. A 5-fluorouracil and leucovorin-containing regimen (off-label);
 - c. A capecitabine-containing regimen (off-label);
5. Prescribed in combination with irinotecan or FOLFIRI;
6. Request meets one of the following (a or b):*
 - a. Dose does not exceed 4 mg/kg every 2 weeks;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 12 months

CLINICAL POLICY

Ziv-Aflibercept

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255.
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53.

II. Continued Therapy

A. Colorectal Cancer (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connections benefit, or documentation supports that member is currently receiving Zaltrap for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed 4 mg/kg every 2 weeks;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255.
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – LA.PMN.53.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CapeOX: capecitabine and oxaliplatin

CRC: colorectal cancer

FDA: Food and Drug Administration

FOLFIRI: fluorouracil, leucovorin, irinotecan

FOLFOX: fluorouracil, leucovorin, oxaliplatin

VEGF: vascular endothelial growth factor

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

CLINICAL POLICY
Ziv-Aflibercept

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Modified FOLFOX 6	Day 1: oxaliplatin 85 mg/m ² IV Day 1: Folinic acid 400 mg/m ² IV Days 1–3: 5-FU 400 mg/m ² IV bolus on day 1, then 1,200 mg/m ² /day × 2 days (total 2,400 mg/m ² over 46–48 hours) IV continuous infusion. Repeat cycle every 2 weeks.	See dosing regimen
CapeOX	Day 1: Oxaliplatin 130 mg/m ² IV Days 1–14: Capecitabine 1,000 mg/m ² PO BID. Repeat cycle every 3 weeks.	See dosing regimen
FOLFIRI	Day 1: Irinotecan 180 mg/m ² IV Day 1: Leucovorin 400 mg/m ² IV Day 1: Fluorouracil 400 mg/m ² IV followed by 2400 mg/m ² continuous IV over 46 hours Repeat cycle every 14 days.	See dosing regimen
5-fluorouracil and leucovorin	Roswell Park regimen: Leucovorin 500 mg/m ² IV followed by 5-FU 500 mg/m ² IV bolus one hour after start of leucovorin on days 1, 8, 15, 22, 29, 36. Repeat every 8 weeks. Biweekly regimen: Leucovorin 400 mg/m ² IV on day one followed by 5-FU 400 mg/m ² IV bolus then 1,200 mg/m ² continuous IV. Repeat every 2 weeks. Weekly regimen: Leucovorin 20 mg/m ² IV on day one followed 5-FU 500 mg/m ² IV bolus one hour after start of leucovorin. Alternatively 5-FU 2,600 mg/m ² continuous IV with leucovorin 500 mg/m ² IV. Repeat weekly.	See dosing regimen
capecitabine	850 – 1,250 mg/m ² PO BID on days 1-14. Repeat every 3 weeks.	2,500 mg/m ² /day

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

None reported

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
CRC	4 mg/kg IV over 1 hour every two weeks	4 mg/kg

VI. Product Availability

Single-use vials for injection: 100 mg/4 mL, 200 mg/8 mL

VII. References

1. Zaltrap Prescribing Information. Bridgewater, NJ: Sanofi-Aventis U.S., LLC; May 2025. Available at https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/125418s0541bl.pdf. Accessed August 4, 2026.
2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed August 4, 2026.
3. National Comprehensive Cancer Network. Colon Cancer Version 2.2026. Available at: https://www.nccn.org/professionals/physician_gls/pdf/colon.pdf. Accessed August 4, 2026.
4. National Comprehensive Cancer Network. Rectal Cancer Version 2.2026. Available at: https://www.nccn.org/professionals/physician_gls/pdf/rectal.pdf. Accessed August 4, 2026.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J9400	Injection, ziv-aflibercept, 1 mg

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Converted corporate to local policy.	06.16.23	10.05.23
Annual review: no significant changes; references reviewed and updated.	04.09.24	07.10.24
No significant changes; references reviewed and updated.	12.03.24	02.24.25
Annual review: extended initial approval duration from 6 to 12 months; references reviewed and updated.	11.20.25	04.08.26
Annual review: no significant changes; references reviewed and updated.	09.21.26	N/A, no material changes

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

CLINICAL POLICY

Ziv-Aflibercept

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2025 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.