

Clinical Policy: Denosumab (Xgeva)

Reference Number: LA.PHAR.58

Effective Date: 04.21

Last Review Date: 06.27.23

Line of Business: Medicaid

Coding Implications
Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Please note: This policy is for medical benefit

Description

Denosumab (Xgeva®) is a receptor activator of nuclear factor kappa-B ligand inhibitor.

FDA Approved Indication(s)

Xgeva is indicated:

- <u>Multiple myeloma (MM) and solid tumors</u>: For the prevention of skeletal-related events in patients with MM and in patients with bone metastases from solid tumors.
- <u>Giant cell tumor of the bone</u>: For the treatment of adults and skeletally mature adolescents with giant cell tumor of bone that is unresectable or where surgical resection is likely to result in severe morbidity.
- <u>Hypercalcemia of malignancy</u>: For the treatment of hypercalcemia of malignancy refractory to bisphosphonate therapy.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

Index

- I. Initial Approval Criteria
 - **A.** Multiple Myeloma or Solid Tumor (*Xgeva*)
 - **B.** Giant Cell Tumor of Bone (*Xgeva*)
 - **C.** Hypercalcemia of Malignancy (*Xgeva*)
 - **D.** Systemic Mastocytosis (off-label) (*Xgeva*)
 - **E.** Other diagnoses/indications
- II. Continued Therapy
 - **A.** All Indications in Section I (*Xgeva*)
 - **B.** Other diagnoses/indications
- III. Diagnoses/Indications for which coverage is NOT authorized
- IV. Appendices/General Information
- V. Dosage and Administration
- VI. Product Availability
- VII. References

It is the policy of Louisiana Healthcare Connections that Xgeva is **medically necessary** when the following criteria are met:



I. Initial Approval Criteria

A. Multiple Myeloma or Solid Tumor (must meet all):

- 1. Request is for Xgeva;
- 2. Diagnosis of one of the following (a or b):
 - a. MM, and member is receiving or initiating therapy (e.g., chemotherapy, transplant) for symptomatic disease;
 - b. Bone metastasis secondary to solid tumor (e.g., breast, kidney, lung, prostate, thyroid);
- 3. Prescribed by or in consultation with an oncologist;
- 4. Age \geq 18 years or documentation of closed epiphyses on x-ray;
- 5. For indications other than prostate or breast cancer, member meets one of the following (a or b):
 - a. Failure of zoledronic acid* (Zometa) or pamidronate* at up to maximally indicated doses, unless clinically significant adverse effects are experienced or both are contraindicated (see Appendices B and D); *Prior authorization may be required.
 - b. Request is for Stage IV or metastatic cancer or associated conditions. Exception if "clinically equivalent therapy, contains identical active ingredient(s), and proven to have same efficacy";
- 6. Xgeva is not prescribed concurrently with Prolia;
- 7. Dose does not exceed 120 mg every 4 weeks.

Approval duration: 6 months

B. Giant Cell Tumor of Bone (must meet all):

- 1. Request is for Xgeva;
- 2. Diagnosis of giant cell tumor of bone that is characterized as one of the following (a or b):
 - a. Metastatic or unresectable disease;
 - b. Localized disease, and Xgeva is prescribed as a single agent or in combination with interferon alfa or radiation therapy;
- 3. Prescribed by or in consultation with an oncologist;
- 4. Age \geq 18 years or documentation of closed epiphyses on x-ray;
- 5. Xgeva is not prescribed concurrently with Prolia;
- 6. Dose does not exceed 120 mg every 4 weeks plus 120 mg on days 8 and 15 of first month of therapy.

Approval duration: 6 months

C. Hypercalcemia of Malignancy (must meet all):

- 1. Request is for Xgeva;
- 2. Diagnosis of hypercalcemia of malignancy:
- 3. Prescribed by or in consultation with an oncologist;
- 4. Age \geq 18 years or documentation of closed epiphyses on x-ray;
- 5. Albumin-corrected calcium > 12.5 mg/dL despite IV bisphosphonate therapy in the last 30 days (*see Appendix B*);
 - *Prior authorization may be required.
- 6. Xgeva is not prescribed concurrently with Prolia;



7. Dose does not exceed 120 mg every 4 weeks plus 120 mg on days 8 and 15 of first month of therapy.

Approval duration: 6 months

D. Systemic Mastocytosis (off-label) (must meet all):

- 1. Request is for Xgeva;
- 2. Diagnosis of systemic mastocytosis;
- 3. Member has osteopenia or osteoporosis with bone pain:
- 4. Prescribed by or in consultation with an oncologist;
- 5. Age \geq 18 years or documentation of closed epiphyses on x-ray;
- 6. Member meets one of the following (a or b):
 - a. Failure of zoledronic acid* (Zometa) or pamidronate* at up to maximally indicated doses unless clinically significant adverse effects are experienced or both are contraindicated (see Appendices B and D); *Prior authorization may be required.
 - b. Request is for Stage IV or metastatic cancer or associated conditions. Exception if "clinically equivalent therapy, contains identical active ingredient(s), and proven to have same efficacy";
- 7. Xgeva is not prescribed concurrently with Prolia;
- 8. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).*

*Prescribed regimen must be FDA-approved or recommended by NCCN.

Approval duration: 6 months

E. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

II. Continued Therapy

A. All Indications in Section I (must meet all):

- 1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Louisiana Healthcare Connections benefit or member has previously met initial approval criteria;
 - b. Documentation supports that member is currently receiving Xgeva for a covered cancer-related indication and has received this medication for at least 30 days;
- 2. Member is responding positively to therapy;
- 3. If request is for a dose increase, new dose does not exceed 120 mg every 4 weeks or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).*

*Prescribed regimen must be FDA-approved or recommended by NCCN.

Approval duration: 12 months



B. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
- 2. If the requested use (e.g. diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – LA.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

ADT: androgen deprivation therapy GIO: glucocorticoid-induced osteoporosis

BMD: bone mineral density MM: multiple myeloma

FDA: Food and Drug Administration PMO: postmenopausal osteoporosis

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval

criteria. The drugs listed here may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
IV bisphosphonates		
ibandronate (Boniva)	Treatment: PMO	Varies
	Hypercalcemia of malignancy	See prescribing
zoledronic acid (Reclast®;	Reclast:	information and
Zometa)	Treatment/prevention: PMO, GIO	compendia for
	Treatment: male osteoporosis	dosing.
	Treatment: Paget disease	
	Zometa:	
	MM	
	Bone metastasis from solid tumors	
	Hypercalcemia of malignancy	
	Systemic mastocytosis (off-label)	
	Fracture prevention - breast/prostate	
	cancer (off-label)	
pamidronate	MM	
	Bone metastasis from breast cancer	
	Hypercalcemia of malignancy	
	Systemic mastocytosis (off-label)	
	Fracture prevention – breast/prostate	
	cancer (off-label)	



Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s):
 - o Xgeva: hypocalcemia, known clinically significant hypersensitivity to Xgeva
- Boxed warning(s): none reported

Appendix D: IV/PO Bisphosphonates: Examples of Contraindications and Adverse Effects

Appendix D: IV/PO Bisphosphonates: Examples of Contraindications and Adverse Effects				
Bisphosphonates	Oral	IV		
	Formulations	Formulations		
Contraindications				
Hypocalcemia	X	X		
Increased risk of aspiration	X	-		
Hypersensitivity to product component	X	X		
Inability to stand/sit upright for at least 30	X	-		
minutes				
Creatinine clearance < 35 mL/min or evidence of	-	X		
acute renal impairment				
Esophagus abnormalities which delay emptying	X	-		
such as stricture or achalasia				
Clinically significant warnings or adverse side eff	ects	_		
Pregnancy	X	X		
Eye inflammation	X	X		
Acute renal failure	X	X		
Osteonecrosis of the jaw	X	X		
Atypical femoral shaft fracture	X	X		
Drug interactions (product-specific)	X	X		
Severe or incapacitating musculoskeletal pain	X	X		

V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum
D 1	101	120 99	Dose
Denosumab	MM	120 mg SC once every	120
(Xgeva)	Solid tumor - bone metastasis	4 weeks	mg/dose
	Giant cell tumor of bone	120 mg SC every 4	120
	Hypercalcemia of malignancy	weeks plus 120 mg on	mg/dose
		Days 8 and 15 of first	
		month of therapy	

VI. Product Availability

Drug Name	Availability
Denosumab (Xgeva)	Injection (single-use vial): 120 mg/1.7 mL (70 mg/mL)



VII. References

- 1. Xgeva Prescribing Information. Thousand Oaks, CA: Amgen Inc.; June 2020. Available at: http://www.xgeva.com. Accessed October 28, 2022.
- 2. Clinical Pharmacology [database online]. Tampa, FL: Elsevier. 2022. URL: www.clinicalkeys.com/pharmacology.

Oncology

- 3. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at www.nccn.org. Accessed October 28, 2022.
- 4. National Comprehensive Cancer Network. Multiple Myeloma Version 1.2023. Available at: https://www.nccn.org/professionals/physician_gls/pdf/myeloma.pdf. Accessed October 28, 2022.
- 5. National Comprehensive Cancer Network. Bone Cancer Version 2.2023. Available at: https://www.nccn.org/professionals/physician_gls/pdf/bone.pdf. Accessed October 28, 2022.
- 6. National Comprehensive Cancer Network. Breast Cancer Version 4.2022. Available at: https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf. Accessed October 28, 2022.
- 7. National Comprehensive Cancer Network. Prostate Cancer Version 1.2023. Available at: https://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf. Accessed October 28, 2022.
- 8. National Comprehensive Cancer Network. Non-Small Cell Lung Cancer Version 5.2022. Available at: https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf. Accessed October 28, 2022.
- 9. National Comprehensive Cancer Network. Kidney Cancer Version 3.2023. Available at: https://www.nccn.org/professionals/physician_gls/pdf/kidney.pdf. Accessed October 28, 2022.
- 10. National Comprehensive Cancer Network. Systemic Mastocytosis Version 2.2022. Available at: https://www.nccn.org/professionals/physician_gls/pdf/mastocytosis.pdf. Accessed October 28, 2022.
- 11. National Comprehensive Cancer Network. Thyroid Carcinoma Version 2.2022. Available at: https://www.nccn.org/professionals/physician_gls/pdf/thyroid.pdf. Accessed October 28, 2022.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS	Description
Codes	
J0897	Injection, denosumab, 1 mg

Reviews, Revisions, and Approvals	Date	LDH
		Approval Date
Converted corporate to local policy.	01.21	04.21



Reviews, Revisions, and Approvals	Date	LDH Approval Date
Removed Prolia criteria. LDH Prolia criteria utilized for Physician	04.22	07.01.22
Administered Medication Prior Authorizations. For multiple myeloma		
or solid tumor and systemic mastocytosis: allowed bypassing of		
redirection of step therapy in Stage IV or metastatic cancer settings.		
Template changes applied to other diagnoses/indications and continued	06.27.23	
therapy section. References reviewed and updated.		
Added blurb this policy is for medical benefit only.		
Minor grammatical and formatting edits.		
Updated maximum dosage for MM.		

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.



Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2023 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.