

**Clinical Policy: Taliglucerase Alfa (Elelyso)****Reference Number: LA.PHAR.157****Effective Date:****Last Review Date: 01.21****Line of Business: Medicaid****Coding Implications  
Revision Log**

**See Important Reminder at the end of this policy for important regulatory and legal information.**

**Description**

Taliglucerase alfa (Elelyso<sup>®</sup>) is a hydrolytic lysosomal glucocerebroside-specific enzyme.

**FDA Approved Indication(s)**

Elelyso is indicated for the treatment of patients 4 years and older with a confirmed diagnosis of type 1 Gaucher disease (GD1).

**Policy/Criteria**

*Prior authorization is required. Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of Louisiana Healthcare Connections that Elelyso is medically necessary when the following criteria are met:

**I. Initial Approval Criteria****A. Gaucher Disease (must meet all):**

1. Diagnosis of Type 1 (GD1) or Type 3 Gaucher disease (GD3) confirmed by one of the following (a or b):
  - a. Enzyme assay demonstrating a deficiency of beta-glucocerebrosidase (glucosidase) activity;
  - b. DNA testing;
2. Age  $\geq$  4 years;
3. Member is symptomatic (e.g., anemia, thrombocytopenia, bone disease, hepatomegaly, splenomegaly);
4. Elelyso is not prescribed concurrently with Vpriv<sup>®</sup> (velaglucerase alfa) or Cerezyme<sup>®</sup> (imiglucerase).

Approval duration:

Medicaid – 6 months

**B. Other diagnoses/indications**

1. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): LA.PMN.53 for Medicaid.

**II. Continued Therapy****A. Gaucher Disease (must meet all):**

## CLINICAL POLICY

### Taliglucerase Alfa

1. Currently receiving medication via Louisiana Healthcare Connections benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy as evidenced by increased or stabilized platelet count or hemoglobin, reduced or stabilized spleen or liver volume, decreased bone pain;
3. Elelyso is not prescribed concurrently with Vpriv® (velaglucerase alfa) or Cerezyme® (imiglucerase).

Approval duration:

Medicaid – 12 months

**B.** Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Louisiana Healthcare Connections benefit and documentation supports positive response to therapy.  
Approval duration: Duration of request or 6 months (whichever is less); or
2. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): LA.PMN.53 for Medicaid.

**III.** Diagnoses/Indications for which coverage is NOT authorized:

- A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – LA.PMN.53 for Medicaid, or evidence of coverage documents.

**IV.** Appendices/General Information

*Appendix A: Abbreviation/Acronym Key*

ERT: enzyme replacement therapy

GD1: type 1 Gaucher disease

FDA: Food and Drug Administration

GD3: type 3 Gaucher disease

*Appendix B: Therapeutic Alternatives*

Not applicable

*Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s): none reported.
- Boxed warning(s): none reported.

*Appendix D: General Information*

- Measures of Therapeutic Response: GD1 is a heterogeneous disorder which involves the visceral organs, bone marrow, and bone in almost all affected patients. Common conditions resulting from GD1 include anemia, thrombocytopenia, hepatomegaly, splenomegaly, and bone disease. Therefore, hemoglobin level, platelet count, liver volume, spleen volume, and bone pain are clinical parameters that can indicate therapeutic response to GD1 therapies. In some clinical trials, stability has been defined as the following thresholds of change from baseline: hemoglobin level < 1.5 g/dL

## CLINICAL POLICY

### Taliglucerase Alfa

decrease, platelet count < 25% decrease, liver volume < 20% increase, and spleen volume < 25% increase.

- Enzyme replacement therapy such as Cerezyme may have beneficial palliative effects in Type 2 disease, but does not alter the outcome and is not generally used.
- According to the European consensus guidelines revised recommendations on the management of neuronopathic Gaucher disease by Vellodi et al: (1) there is clear evidence in most patients that enzyme replacement therapy (ERT) ameliorates systemic involvement in non-neuronopathic (Type 1) as well as chronic neuronopathic Gaucher disease (Type 3), enhancing quality of life; (2) There is no evidence that ERT has reversed, stabilized or slowed the progression of neurological involvement; (3) In patients with established acute neuronopathic Gaucher disease (Type 2), enzyme replacement therapy has had little effect on the progressively downhill course. It has merely resulted in prolongation of pain and suffering.
- There is currently insufficient clinical evidence that supports the combination use of enzyme replacement therapy with Zavesca® (miglustat) or Cerdelga® (eliglustat), or concurrent use of two or more enzyme replacement therapies at once.

#### V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Gaucher disease	Treatment-naïve patients: 60 units/kg IV every other week  Patients switching from imiglucerase: Begin at the same unit/kg dose as the patient's previous imiglucerase dose; administer every other week	Individualized (adjust based on achievement and maintenance of therapeutic goals)

#### VI. Product Availability

Single-use vial: 200 units

#### VII. References

1. Elelyso Prescribing Information. New York, NY: Pfizer, Inc.; November 2020. Available at <http://www.elelyso.com>. Accessed November 24, 2020.
2. Charrow J, Andersson HC, Kaplan P. Enzyme replacement therapy and monitoring for children with type 1 Gaucher disease: consensus recommendations. J Pediatr. 2004; 144: 112-20.
3. Hollak, CEM, Weinreb NJ. The attenuated/late onset lysosomal storage disorders: therapeutic goals and indications for enzyme replacement treatment in Gaucher and Fabry disease. Best Pract Res Clin Endocrinol Metab. 2015; 29: 205-218.
4. Pastores GM, Weinreb NJ, Aerts H, et al. Therapeutic goals in the treatment of Gaucher disease. Semin Hematol. 2004; 41(suppl 5): 4-14.
5. Andersson HC, Charrow J, Kaplan P, et al. Individualization of long-term enzyme replacement therapy for Gaucher disease. Genet Med. 2005; 7(2): 105-110.
6. Altarescu G, Hill S, Wiggs E, et al. The efficacy of enzyme replacement therapy in patients with chronic neuronopathic Gaucher's disease. J Pediatr. 2001;138:539-547.

## CLINICAL POLICY

### Taliglucerase Alfa

7. Vellodi A, Tylki-Szymanska A, Davies E, et al. Management of neuronopathic Gaucher disease: Revised recommendations. *J Inherit Metab Dis.* 2009;32:660-664.

#### Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J3060	Injection, taliglucerase alfa, 10 units

Reviews, Revisions, and Approvals	Date
Converted corporate to local policy	01.21

#### Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between

## **CLINICAL POLICY**

### Taliglucerase Alfa

the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2020 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.