

Clinical Policy: Aflibercept (Eylea)

Reference Number: LA.PHAR.184

Effective Date: 09.21

Last Review Date: 04.22

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Aflibercept (Eylea®) is a vascular endothelial growth factor (VEGF) inhibitor.

FDA Approved Indication(s)

Eylea is indicated for the treatment of patients with:

- Neovascular (wet) age-related macular degeneration (AMD)
- Macular edema following retinal vein occlusion (RVO)
- Diabetic macular edema (DME)
- Diabetic retinopathy (DR)

Policy/Criteria

Prior authorization is required. Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections that Eylea is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Ophthalmic Disease (must meet all):

1. Diagnosis of one of the following (a, b, c, or d):
 - a. Neovascular (wet) AMD;
 - b. Macular edema following RVO;
 - c. DME;
 - d. DR;
2. Prescribed by or in consultation with an ophthalmologist;
3. Age \geq 18 years;
4. For all indications, except for DME in members with baseline best corrected visual acuity (BCVA) worse than 20/50: Failure of bevacizumab intravitreal solution, unless contraindicated or clinically significant adverse effects are experienced;
**Prior authorization may be required for bevacizumab intravitreal solution. Requests for IV formulations of Avastin, Mvasi, and Zirabev will not be approved.*
5. Dose does not exceed:
 - a. AMD: 2 mg (1 vial) every 4 weeks for the first 3 months, then every 8 weeks thereafter;
 - b. DME and DR: 2 mg (1 vial) every 4 weeks for the first 5 injections, then every 8 weeks thereafter;
 - c. RVO: 2 mg (1 vial) every 4 weeks.

Approval duration: 6 months

B. Other diagnoses/indications

1. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): LA.PMN.53 for Medicaid.

II. Continued Therapy

A. Ophthalmic Disease (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connections benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy as evidenced by one of the following (a, b, c, or d):
 - a. Detained neovascularization;
 - b. Improvement/stabilization in visual acuity;
 - c. Maintenance of corrected visual acuity from prior treatment;
 - d. Supportive findings from optical coherence tomography or fluorescein angiography;
3. If request is for a dose increase, new dose does not exceed:
 - a. DME and DR: 2 mg (1 vial) every 8 weeks;
 - b. RVO: 2 mg (1 vial) every 4 weeks;
 - c. AMD: One of the following (i or ii):
 - i. Dose does not exceed 2 mg (1 vial) every 8 weeks;
 - ii. Member meets both of the following (a and b):
 - a) Documentation supports evidence of continued disease activity;
 - b) New dose does not exceed 2 mg (1 vial) every 4 weeks.

Approval duration: 6 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Louisiana Healthcare Connections benefit and documentation supports positive response to therapy.
Approval duration: Duration of request or 6 months (whichever is less); or
2. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): LA.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – LA.PMN.53 for Medicaid or or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

AMD: age-related macular degeneration

BCVA: best corrected visual acuity

DME: diabetic macular edema

DR: diabetic retinopathy

CLINICAL POLICY

Aflibercept

FDA: Food and Drug Administration
RVO: retinal vein occlusion

VEGF: vascular endothelial growth factor

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Bevacizumab (Avastin®)	Neovascular (wet) AMD: 1.25 to 2.5 mg administered by intravitreal injection every 4 weeks.	2.5 mg/month
	Macular edema secondary to RVO: 1 mg to 2.5 mg administered by intravitreal injection every 4 weeks	2.5 mg/month
	DR: 1.25 mg administered by intravitreal injection every 6 weeks	1.25 mg/6 weeks
	DME: 1.25 mg administered by intravitreal injection every 6 weeks	1.25 mg/6 weeks

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s):
 - Ocular or periocular infection
 - Active intraocular inflammation
 - Hypersensitivity
- Boxed warning(s): none reported

Appendix D: General Information

- In the VEGF Trap-Eye: Investigation of Efficacy and Safety in Wet Age-Related Macular Degeneration (VIEW)-1 trial, the difference in the number of patients who lost fewer than 15 letters at 52 weeks between Eylea every 8 weeks compared to Lucentis was 0.6% (95.1% CI -0.32, 4.4). In terms of the number of patients who gained at least 15 letters, the mean difference between Eylea every 8 weeks was 6.6% (95.1% CI -1.0, 14.1). There were no adverse events that were found to be significant from the Lucentis arm.
- In a trial comparing Eylea, Avastin and Lucentis, the Diabetic Retinopathy Clinical Research Network found in patients with diabetic macular edema that when the initial visual-acuity letter score was 78 to 69 (equivalent to approximately 20/32 to 20/40) (51% of participants), the mean improvement was 8.0 with Eylea, 7.5 with Avastin, and 8.3 with Lucentis ($p > 0.50$ for each pair wise comparison). When the initial letter score was less than 69 (approximately 20/50 or worse), the mean improvement was 18.9 with Eylea,

CLINICAL POLICY

Aflibercept

11.8 with Avastin, and 14.2 with Lucentis ($p < 0.001$ for Eylea vs. Avastin, $p = 0.003$ for Eylea vs. Lucentis, and $p = 0.21$ for Lucentis vs. Avastin).

- In clinical trials for the treatment of AMD, DME, and DR, additional efficacy was not demonstrated in most patients when Eylea was dosed every 4 weeks as a maintenance dose, compared to every 8 weeks. Maintenance dosing at every 8 weeks should be attempted before increasing the intravitreal injection frequency to every 4 weeks.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
AMD	2 mg (1 vial) administered by intravitreal injection once a month for 3 months then 2 mg every 2 months <i>Although Eylea may be dosed as frequently as 2 mg every 4 weeks (monthly), additional efficacy was not demonstrated in most patients when Eylea was dosed every 4 weeks compared to every 8 weeks. Some patients may need every 4 week (monthly) dosing after the first 12 weeks (3 months).</i>	2 mg/month
Macular edema following RVO	2 mg (1 vial) administered by intravitreal injection once every 4 weeks (monthly)	2 mg/month
DME, DR	2 mg (1 vial) administered by intravitreal injection once a month for the first 5 injections, followed by 2 mg via intravitreal injection once every 2 months <i>Although Eylea may be dosed as frequently as 2 mg every 4 weeks (monthly), additional efficacy was not demonstrated in most patients when Eylea was dosed every 4 weeks compared to every 8 weeks. Some patients may need every 4 week (monthly) dosing after the first 20 weeks (5 months).</i>	2 mg/month

VI. Product Availability

Single-dose vial and pre-filled syringe: 2 mg/0.05 mL solution

VII. References

1. Eylea Prescribing Information. Tarrytown, NY: Regeneron Pharmaceuticals, Inc.; March 2021. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125387s069lbl.pdf. Accessed November 9, 2021.
2. American Academy of Ophthalmology Retina/Vitreous Panel. Preferred Practice Pattern® Guidelines. Age-Related Macular Degeneration. San Francisco, CA: American Academy

CLINICAL POLICY

Aflibercept

of Ophthalmology; September 2019. Available at: www.aao.org/ppp. Accessed November 9, 2021.

3. American Academy of Ophthalmology Retina/Vitreous Panel. Preferred Practice Pattern® Guidelines. Retinal Vein Occlusions. San Francisco, CA: American Academy of Ophthalmology; September 2019. Available at: www.aao.org/ppp. Accessed November 9, 2021.
4. American Academy of Ophthalmology Retina/Vitreous Panel. Preferred Practice Pattern® Guidelines. Diabetic Retinopathy. San Francisco, CA: American Academy of Ophthalmology; September 2019. Available at: www.aao.org/ppp. Accessed November 9, 2021.
5. Wells JA, Glassman AR, Ayala AR, et al. Aflibercept, bevacizumab, or ranibizumab for diabetic macular edema. N Engl J Med. 2015 Mar 26;372(13):1193-203. Doi: 10.1056/NEJMoa1414264.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J0178	Injection, aflibercept, 1 mg

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Converted corporate to local policy	06.21	09.21
Clarified “best corrected” for visual acuity for redirection to bevacizumab. Converted redirection language from “must use” to “Failure of” bevacizumab intravitreal solution. References reviewed and updated.	04.22	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

CLINICAL POLICY

Aflibercept

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2020 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.