

Clinical Policy: Pertuzumab (Perjeta)

Reference Number: LA.PHAR.227

Effective Date:

Last Review Date: 04.22 Coding Implications
Line of Business: Medicaid Revision Log

See Important Reminder at the end of this policy for important regulatory and legal information.

Description

Pertuzumab (Perjeta®) is a human epidermal growth factor receptor 2 protein (HER2)/neu receptor antagonist.

FDA Approved Indication(s)

Perjeta is indicated for:

- Use in combination with trastuzumab and docetaxel for the treatment of patients with HER2positive metastatic breast cancer who have not received prior anti-HER2 therapy or chemotherapy for metastatic disease.
- Use in combination with trastuzumab and chemotherapy as:
 - Neoadjuvant treatment of patients with HER2-positive, locally advanced, inflammatory, or early stage breast cancer (either greater than 2 cm in diameter or node positive) as part of a complete treatment regimen for early breast cancer;
 - o Adjuvant treatment of patients with HER2-positive early breast cancer at high risk of recurrence.

Policy/Criteria

Prior Authorization is required. Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections that Perjeta is medically necessary when the following criteria are met:

- I. Initial Approval Criteria
 - **A.** Breast Cancer (must meet all):
 - 1. Diagnosis of HER2-positive breast cancer;
 - 2. Prescribed by or in consultation with an oncologist;
 - 3. Age \geq 18 years;
 - 4. Prescribed as combination therapy (see Appendix B);
 - 5. Request meets one of the following (a or b):*
 - a. Initial dose: 840 mg, followed by maintenance dose: 420 mg every 3 weeks;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration: 6 months

- **B.** Additional NCCN Recommended Uses (off-label) (must meet all):
 - 1. Diagnosis of one of the following (a or b):

Pertuzumah



- a. Recurrent HER2-positive salivary gland tumor;
- b. Advanced or metastatic colorectal cancer and disease is all of the following (i, ii, and iii):
 - i. HER2 positive;
 - ii. Wild-type *RAS* (defined as wild-type in both KRAS and NRAS as determined by an FDA-approved test for this use);
 - iii. Wild-type *BRAF*;
- 2. Prescribed by or in consultation with an oncologist;
- 3. Age \geq 18 years;
- 4. For colorectal cancer: No previous use of a HER2 inhibitor therapy (e.g., trastuzumab, Kadcyla[®], Tykerb[®], Perjeta);
- 5. Prescribed in combination with trastuzumab;* **Prior authorization may be required.*
- 6. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).*

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration: 6 months

C. Other diagnoses/indications

 Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): LA.PMN.53 for Medicaid.

II. Continued Therapy

- **A.** All Indications in Section I (must meet all):
 - 1. Currently receiving medication via Louisiana Healthcare Connections benefit, or documentation supports that member is currently receiving Perjeta for a covered indication and has received this medication for at least 30 days;
 - 2. Member is responding positively to therapy;
 - 3. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed 420 mg every 3 weeks;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration: 12 months (total of 18 cycles if neoadjuvant or adjuvant therapy)

- **B.** Other diagnoses/indications (must meet 1 or 2):
 - 1. Currently receiving medication via Louisiana Healthcare Connections benefit and documentation supports positive response to therapy.
 - Approval duration: Duration of request or 6 months (whichever is less); or
 - 2. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): LA.PMN.53 for Medicaid.

Pertuzumab



III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – LA.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

BRAF: v-raf murine sarcoma viral

oncogene homolog B1

FDA: Food and Drug Administration

HER2: human epidermal growth factor

receptor 2

KRAS: Kirsten rat sarcoma 2 viral oncogene homologue

NRAS: neuroblastoma RAS viral oncogene

homologue

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval

criteria. The drugs listed here may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum
Examples of drugs that may be used with Perjeta for breast cancer: • Chemotherapeutic agents: carboplatin, cyclophosphamide, doxorubicin, docetaxel, paclitaxel • HER2-targeted agents: docetaxel (Taxotere®), paclitaxel, Herceptin® (trastuzumab) • Endocrine therapy: tamoxifen; aromatase inhibitors: anastrozole (Arimidex®), letrozole (Femara®), exemestane (Aromasin®).	Regimens are dependent on a variety of factors including menopausal status, treatment/progression history, clinical stage, histology, mutational and receptor status, treatment purpose (e.g., adjuvant and neoadjuvant treatment, treatment for metastatic disease).	Varies Varies

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): Known hypersensitivity to pertuzumab or to any of its excipients
- Boxed warning(s): Left ventricular dysfunction, embryo-fetal toxicity

V. Dosage and Administration

Indication	Dosing Regimen	Maximum
		Dose
Breast	Initial dose of 840 mg IV, followed by maintenance dose of 420	See
cancer	mg IV every 3 weeks	regimens

louisiana healthcare connections

Pertuzumab

Indication	Dosing Regimen	Maximum Dose
	For metastatic disease, Perjeta should be administered as outlined above. For neoadjuvant treatment, Perjeta should be administered for 3-6 cycles. Following surgery, patients should continue to receive Perjeta to complete 1 year of treatment (up to 18 cycles) For adjuvant treatment, Perjeta should be administered for a total of 1 year (up to 18 cycles) or until disease recurrence or unmanageable toxicity.	

VI. Product Availability

Single-dose vial for injection: 420 mg/14 mL

VII. References

- 1. Perjeta Prescribing Information. South San Francisco, CA: Genentech, Inc.; February 2021. Available at https://www.gene.com/download/pdf/perjeta_prescribing.pdf. Accessed February 15, 2022.
- 2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at www.nccn.org. Accessed February 15, 2022.
- 3. National Comprehensive Cancer Network Guidelines. Breast Cancer Version 2.2022. Available at www.nccn.org. Accessed February 15, 2022.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J9306	Injection, pertuzumab, 1 mg

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Converted corporate to local policy.	04.22	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical

Pertuzumab



policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2020 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana



Pertuzumab

Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.