

Clinical Policy: Amivantamab-vmjw (Rybrevant)

Reference Number: LA.PHAR.544

Effective Date:

Last Review Date: 05.01.23

Line of Business: Medicaid

[Coding Implications](#)
[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

****Please note: This policy is for medical benefit****

Description

Amivantamab-vmjw (Rybrevant™) is a bispecific epidermal growth factor (EGF) receptor-directed and MET receptor-directed antibody.

FDA Approved Indication(s)

Rybrevant is indicated for the treatment of adult patients with locally advanced or metastatic non-small cell lung cancer (NSCLC) with epidermal growth factor receptor (EGFR) exon 20 insertion mutations, as detected by an FDA-approved test, whose disease has progressed on or after platinum-based chemotherapy.

This indication is approved under accelerated approval based on overall response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in the confirmatory trials.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections® that Rybrevant is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Non-Small Cell Lung Cancer (must meet all):

1. Diagnosis of recurrent, locally advanced or metastatic NSCLC;
2. Prescribed by or in consultation with an oncologist;
3. Age \geq 18 years;
4. Disease is positive for epidermal growth factor receptor (EGFR) exon 20 insertion mutations;
5. Member has progressed on or after platinum-based therapy;
6. Request meets one of the following (a or b):*
 - a. Dose does not exceed the appropriate weight-based dose (i or ii) per week for 4 weeks, then every 2 weeks thereafter (*see section V for dosing regimen*):
 - i. Body weight < 80 kg: 1,050 mg (3 vials);
 - ii. Body weight \geq 80 kg: 1,400 mg (4 vials);

CLINICAL POLICY

Amivantamab-vmjw

- b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

II. Continued Therapy

A. Non-Small Cell Lung Cancer (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connections benefit, or documentation supports that member is currently receiving Rybrevant for NSCLC and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New Dose does not exceed the appropriate weight-based dose (i or ii) every 2 weeks (*see section V for dosing regimen*):
 - i. Body weight < 80 kg: 1,050 mg (3 vials);
 - ii. Body weight ≥ 80 kg: 1,400 mg (4 vials);
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – LA.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

CLINICAL POLICY

Amivantamab-vmjw

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration

MET: mesenchymal-epithelial transition

EGFR: epidermal growth factor receptor

TKI: tyrosine kinase inhibitor

NSCLC: non-small cell lung cancer

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Platinum-based chemotherapy (e.g., cisplatin, carboplatin)	Varies	Varies

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

None reported

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
NSCLC	<p>Weight-based dose IV weekly for 4 weeks, with the initial dose as a split infusion in Week 1 on Day 1 and Day 2, then every 2 weeks thereafter:</p> <p>Week 1, day 1:</p> <ul style="list-style-type: none"> Body weight < 80 kg: 350 mg (1 vial) Body weight ≥ 80 kg: 350 mg (1 vial) <p>Week 1, day 2:</p> <ul style="list-style-type: none"> Body weight < 80 kg: 700 mg (2 vials) Body weight ≥ 80 kg: 1050 mg (3 vials) <p>Week 2 and thereafter:</p> <ul style="list-style-type: none"> Body weight < 80 kg: 1,050 mg (3 vials) Body weight ≥ 80 kg: 1,400 mg (4 vials) 	See regimen

VI. Product Availability

Solution for injection in a single-dose vial: 350 mg/7 mL (50 mg/mL)

VII. References

1. Rybrevant Prescribing Information. Horsham, PA: Janssen Biotech, Inc.; December 2021. Available at: <https://www.Rybrevant.com/>. Accessed May 3, 2022.
2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed May 3, 2022.

CLINICAL POLICY

Amivantamab-vmjw

3. National Comprehensive Cancer Network. Non-Small Cell Lung Cancer. Version 3.2022.
Available at: http://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf. Accessed May 3, 2022.

CLINICAL POLICY

Amivantamab-vmjw

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPSC Codes	Description
J9061	Injection, amivantamab-vmjw, 2 mg

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Policy created	05.01.23	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care and are solely responsible

CLINICAL POLICY

Amivantamab-vmjw

for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2023 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.