

Clinical Policy: Ferric Carboxymaltose (Injectafer)

Reference Number: CP.PHAR.234

Effective Date: 06.01.16

Last Review Date: 02.19

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Ferric carboxymaltose (Injectafer[®]) injection is an iron replacement product.

FDA Approved Indication(s)

Injectafer is indicated for treatment of iron deficiency anemia (IDA) in adult patients who have:

- Intolerance to oral iron or have had unsatisfactory response to oral iron; or
- Nondialysis-dependent chronic kidney disease (CKD).

Policy/Criteria

Provider must submit documentation (including such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Injectafer is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Iron Deficiency Anemia with Chronic Kidney Disease (must meet all):

1. Diagnosis of IDA and CKD;
2. IDA is confirmed by either of the following:
 - a. Transferrin saturation (TSAT) \leq 30%;
 - b. Serum ferritin \leq 500 ng/mL ;
3. If CKD does not require hemodialysis or peritoneal dialysis, oral iron therapy is not optimal due to any of the following:
 - a. TSAT $<$ 12%;
 - b. Hgb $<$ 7 g/dL;
 - c. Symptomatic anemia;
 - d. Severe or ongoing blood loss;
 - e. Oral iron intolerance;
 - f. Unable to achieve therapeutic targets with oral iron;
 - g. Co-existing condition that may be refractory to oral iron therapy;
4. Dose does not exceed 750 mg elemental iron per infusion/injection.

Approval duration: 3 months

B. Iron Deficiency Anemia without Chronic Kidney Disease (must meet all):

1. Diagnosis of IDA confirmed by any of the following:
 - a. Serum ferritin $<$ 15 ng/mL or $<$ 30 ng/mL if pregnant;
 - b. Serum ferritin \leq 41 ng/mL and Hgb $<$ 12 g/dL (women)/ $<$ 13 g/dL (men);

CLINICAL POLICY
Ferric Carboxymaltose

- c. TSAT < 20%;
- d. Absence of stainable iron in bone marrow;
- e. Increased soluble transferrin receptor (sTfR) or sTfR-ferritin index;
- f. Increased erythrocyte protoporphyrin level;
2. Oral iron therapy is not optimal due to any of the following:
 - a. TSAT < 12%;
 - b. Hgb < 7 g/dL;
 - c. Symptomatic anemia;
 - d. Severe or ongoing blood loss;
 - e. Oral iron intolerance;
 - f. Unable to achieve therapeutic targets with oral iron;
 - g. Co-existing condition that may be refractory to oral iron therapy;
3. At the time of the request, member does not have CKD;
4. Dose does not exceed 750 mg elemental iron per infusion/injection.

Approval duration 3 months**C. Other diagnoses/indications:**

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid.

II. Continued Approval Criteria**A. Iron Deficiency Anemia with Chronic Kidney Disease (must meet all):**

1. Currently receiving the medication via Centene benefit or member has previously met all initial approval criteria;
2. Documentation of one of the following laboratory results measured since the last IV iron administration:
 - a. TSAT \leq 30%;
 - b. Serum ferritin \leq 500 ng/mL;
3. If request is for a dose increase, new dose not exceed 750 mg elemental iron per infusion/injection.

Approval duration 3 months**B. Iron Deficiency Anemia without Chronic Kidney Disease (must meet all):**

1. Currently receiving the medication via Centene benefit or member has previously met all initial approval criteria;
2. Documentation of one of the following laboratory results measured since the last IV iron administration:
 - a. Serum ferritin < 15 ng/mL or < 30 ng/mL if pregnant;
 - b. Serum ferritin \leq 41 ng/mL and Hb < 12 g/dL (women)/< 13 g/dL (men);
 - c. TSAT < 20%;
 - d. Absence of stainable iron in bone marrow;
 - e. Increased sTfR or sTfR-ferritin index;
 - f. Increased erythrocyte protoporphyrin level;
3. At the time of the request, member does not have CKD;

CLINICAL POLICY

Ferric Carboxymaltose

- If request is for a dose increase, new dose does not exceed 750 mg elemental iron per infusion/injection.

Approval duration 3 months

C. Other diagnoses/indications (must meet 1 or 2):

- Currently receiving medication via Centene benefit and documentation supports positive response to therapy.
Approval duration: Duration of request or 6 months (whichever is less); or
- Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

| | |
|---------------------------------------|------------------------------------|
| CKD: chronic kidney disease | IDA: iron deficiency anemia |
| ESA: erythropoiesis stimulating agent | TSAT: transferrin saturation |
| Hb: hemoglobin | sTfR: soluble transferrin receptor |

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

| Drug Name | Dosing Regimen | Dose Limit/Maximum Dose |
|--|----------------|-------------------------|
| Examples of OTC Oral Iron Formulations* | | |
| Ferrous fumarate (Ferretts, Ferrimin 150, Hemocyte) | Varies | |
| Ferrous gluconate (Ferlate) | | |
| Ferrous sulfate (BProtected Pedia Iron, Fer-In-Sol, FeroSul, FerronSul, Iron Supplement Childrens, Slow Fe, Slow Iron) | | |
| Polysaccharide-iron complex (EZFE 200, Ferrex 150, Ferrix x-150, Myferon 150, NovaFerrum 125, NovaFerrum 50, NovaFerrum Pediatric Drops, Nu-Iron, Poly-Iron 150) | | |

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

**Oral formulations include elixirs, liquids, solutions, syrups, capsules, and tablets - including delayed/extended-release tablets.*

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): Hypersensitivity to Injectafer or any of its inactive components.
- Boxed warning(s): None reported.

CLINICAL POLICY
Ferric Carboxymaltose

V. Dosage and Administration

| Indication | Dosing Regimen | Maximum Dose |
|---|---|--|
| IDA with or without non-dialysis dependent CKD (adults) | <p>≥ 50kg (110lb): two 750 mg doses separated by at least 7 days for a cumulative dose of 1500 mg per course.</p> <p>< 50kg (110lb): two doses separated by at least 7 days as 15 mg/kg body weight.</p> | <p>750 mg per dose</p> <p>-Treatment course: up to 1500 mg</p> <p>-Treatment may be repeated</p> |

VI. Product Availability

Intravenous solution: 750 mg/15 mL (15 mL)

VII. References

1. Injectafer prescribing information. Shirley, NY: American Regent, Inc.; April 2018. Available from <https://www.accessdata.fda.gov>. Accessed November 06, 2018.
2. KDIGO 2012 clinical practice guideline for evaluation and management of chronic kidney disease. *Kidney International Supplements*. January 2013; 3(1): 1-136.
3. KDIGO 2012 clinical practice guideline for anemia in chronic kidney disease. *Kidney International Supplements*. August 2012; 2(4): 279-331.
4. Camaschella C. Iron-Deficiency Anemia. *N Engl J Med*. 2015; 372: 1832-43. DOI: 10.1056/NEJMra1401038.
5. Short MW, Domagalski JE. Iron Deficiency Anemia: Evaluation and Management. *Am Fam Physician*. 2013; 87(2): 98-104. <http://www.aafp.org/afp/2013/0115/p98.pdf>
6. Oral iron monographs. In: UpToDate (Lexicomp), Waltham, MA: Walters Kluwer Health; 2018. Available at www.uptodate.com. Accessed November 06, 2018.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

| HCPCS Codes | Description |
|-------------|--|
| J1439 | Injection, ferric carboxymaltose, 1 mg |

| Reviews, Revisions, and Approvals | Date | P&T Approval Date |
|--|-------|-------------------|
| Policy developed | 01.16 | 03.16 |
| Labeled and off-labeled use, and diagnostic/follow-up tests, are edited for consistency across ferumoxytol, ferric gluconate, iron sucrose, and ferric carboxymaltose policies, and are made broad enough to capture use in adults, children and pregnancy. The criteria also encompass iron maintenance and replenishment. Diagnostic hemoglobin for anemia in men changed from 13.5 to | 02.17 | 03.17 |

CLINICAL POLICY
Ferric Carboxymaltose

| Reviews, Revisions, and Approvals | Date | P&T Approval Date |
|--|----------|-------------------|
| 13 based on WHO criteria. Age and dose are removed. Hypersensitivity removed as a contraindication. | | |
| 1Q18 annual review: - No significant changes - Converted to the new template - Dosing added - References reviewed and updated. | 12.01.17 | 02.18 |
| 1Q 2019 annual review; under IDA initial and continuation criteria, a serum ferritin of less than or equal to 500 is edited by deleting the additional requirement of receiving an ESA based on the KDIGO 2012 guidelines which do not include this restriction; under IDA and IDA with CKD continuation criteria, the greater than or equal to 4 week waiting period before retesting after the last IV iron administration is removed per the KDIGO 2012 guidelines which note that only one week need pass before retesting; references reviewed and updated. | 11.13.18 | 02.19 |

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan

CLINICAL POLICY

Ferric Carboxymaltose

retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note: For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.