

## **Clinical Policy: Anakinra (Kineret)**

Reference Number: CP.PHAR.244

Effective Date: 08.16

Last Review Date: 05.18

Line of Business: HIM, Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### **Description**

Anakinra (Kineret<sup>®</sup>) is an interleukin-1 (IL-1) receptor antagonist.

### **FDA Approved Indication(s)**

Kineret is indicated for the treatment of:

- Rheumatoid arthritis (RA): Reduction in signs and symptoms and slowing the progression of structural damage in moderately to severely active RA, in patients 18 years of age or older who have failed 1 or more disease modifying antirheumatic drugs (DMARDs). Kineret can be used alone or in combination with DMARDs other than tumor necrosis factor blocking agents.
- Cryopyrin-associated periodic syndromes (CAPS): Treatment of neonatal-onset multisystem inflammatory disease (NOMID)

### **Policy/Criteria**

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Kineret is **medically necessary** when the following criteria are met:

#### **I. Initial Approval Criteria**

##### **A. Rheumatoid Arthritis** (must meet all):

1. Diagnosis of RA;
2. Prescribed by or in consultation with a rheumatologist;
3. Age  $\geq$  18 years;
4. Member meets one of the following (a or b):
  - a. Failure of a  $\geq$  3 consecutive month trial of methotrexate (MTX) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
  - b. If intolerance or contraindication to MTX (*see Appendix C*), failure of a  $\geq$  3 consecutive month trial of at least ONE conventional DMARD (e.g., sulfasalazine, leflunomide, hydroxychloroquine) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
5. Failure of etanercept (*Enbrel<sup>®</sup> is preferred*) and adalimumab (*Humira<sup>®</sup> is preferred*) each used for  $\geq$  3 consecutive months, unless contraindicated or clinically significant adverse effects are experienced;

*\*Prior authorization is required for etanercept and adalimumab*

6. Dose does not exceed 100 mg per day.

**Approval duration: 6 months**

**B. Cryopyrin-Associated Periodic Syndromes (must meet all):**

1. Diagnosis of NOMID;
2. Prescribed by or in consultation with a rheumatologist;
3. Dose does not exceed 8 mg/kg per day.

**Approval duration: 6 months**

**C. Other diagnoses/indications**

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**II. Continued Therapy**

**A. All Indications in Section I (must meet all):**

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed one of the following (a or b):
  - a. RA: 100 mg per day;
  - b. NOMID: 8mg/kg per day.

**Approval duration: 12 months**

**B. Other diagnoses/indications (must meet 1 or 2):**

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.  
**Approval duration: Duration of request or 6 months (whichever is less);** or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

CAPS: cryopyrin-associated periodic syndromes

DMARD: disease-modifying antirheumatic drug

FDA: Food and Drug Administration  
IL-1: interleukin-1  
MTX: methotrexate

NOMID: neonatal-onset multisystem  
inflammatory disease  
RA: rheumatoid arthritis

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

<b>Drug Name</b>	<b>Dosing Regimen</b>	<b>Dose Limit/ Maximum Dose</b>
azathioprine (Azasan <sup>®</sup> , Imuran <sup>®</sup> )	<b>RA</b> 1 mg/kg/day PO QD or divided BID	2.5 mg/kg/day
Cuprimine <sup>®</sup> (d-penicillamine)	<b>RA*</b> <u>Initial dose:</u> 125 or 250 mg PO QD <u>Maintenance dose:</u> 500 – 750 mg/day PO QD	1,500 mg/day
cyclosporine (Sandimmune <sup>®</sup> , Neoral <sup>®</sup> )	<b>RA</b> 2.5 – 4 mg/kg/day PO divided BID	4 mg/kg/day
hydroxychloroquine (Plaquenil <sup>®</sup> )	<b>RA*</b> <u>Initial dose:</u> 400 – 600 mg/day PO QD <u>Maintenance dose:</u> 200 – 400 mg/day PO QD	600 mg/day
leflunomide (Arava <sup>®</sup> )	<b>RA</b> 100 mg PO QD for 3 days, then 20 mg PO QD	20 mg/day
methotrexate (Rheumatrex <sup>®</sup> )	<b>RA</b> 7.5 mg/week PO, SC, or IM or 2.5 mg PO Q12 hr for 3 doses/week	30 mg/week
Ridaura <sup>®</sup> (auranofin)	<b>RA</b> 6 mg PO QD or 3 mg PO BID	9 mg/day (3 mg TID)
sulfasalazine (Azulfidine <sup>®</sup> )	<b>RA</b> 2 g/day PO in divided doses	3 g/day
Enbrel <sup>®</sup> (etanercept)	<b>RA</b> 25 mg SC twice weekly or 50 mg SC once weekly	50 mg/week
Humira <sup>®</sup> (adalimumab)	<b>RA</b> 40 mg SC every other week (may increase to once weekly)	40 mg/week

*Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.*

*\*Off-label*

*Appendix C: General Information*

- Definition of MTX or DMARD Failure
  - Child-bearing age is not considered a contraindication for use of MTX. Each drug has risks in pregnancy. An educated patient and family planning would allow use of MTX in patients who have no intention of immediate pregnancy.
  - Social use of alcohol is not considered a contraindication for use of MTX. MTX may only be contraindicated if patients choose to drink over 14 units of alcohol per week. However, excessive alcohol drinking can lead to worsening of the condition, so patients who are serious about clinical response to therapy should refrain from excessive alcohol consumption.
- Examples of positive response to therapy may include, but are not limited to:
  - Reduction in joint pain/swelling/tenderness
  - Improvement in ESR/CRP levels
  - Improvements in activities of daily living

**V. Dosage and Administration**

Indication	Dosing Regimen	Maximum Dose
RA	100 mg SC QD	100 mg/day
NOMID	<u>Initial dose:</u> 1 – 2 mg/kg SC QD or divided BID <u>Maintenance dose:</u> 8 mg/kg SC QD or divided BID	8 mg/kg/day

**VI. Product Availability**

Single-use prefilled syringe: 100 mg/0.67 mL

**VII. References**

1. Kineret Prescribing Information. Stockholm, Sweden: Swedish Orphan Biovitrum AB; May 2016. Available at <http://www.kineretrx.com/pdf/Full-Prescribing-Information-English.pdf>. Accessed February 27, 2018.
2. Smolen JS, Landewé R, Breedveld FC, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2013 update. *Ann Rheum Dis*. 2014; 73: 492-509.
3. Singh JA, Furst DE, Bharat A, et al. 2012 update of the 2008 American College of Rheumatology recommendations for the use of disease-modifying antirheumatic drugs and biologic agents in the treatment of rheumatoid arthritis. *Arthritis Care Res*. 2012; 64(5): 625-639.
4. Singh JA, Saag KG, Bridges SL, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Rheumatology* 2016. 68(1):1-26.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy split from CP.PHAR.86.Arthritis Treatments and CP.PHAR.47. CAPS. Removed criteria related to HBV, malignant disease, concomitant use with other biologics, and concurrent	06.16	08.16

Reviews, Revisions, and Approvals	Date	P&T Approval Date
administration of live vaccines; added dosing requirement. RA: changed age requirement to 18 years; modified criteria to require trial of methotrexate, unless contraindicated; added sulfasalazine and hydroxychloroquine as an alternative to MTX if MTX is contraindicated; added requirement for trial and failure of PDL Enbrel and Humira, unless contraindicated. Re-auth: combined into All Indications; added criteria related to dosing per PI and reasons to discontinue. Modified approval duration to 6 months for initial and 12 months for renewal.		
Converted to new template. RA: revised criteria for confirmation of RA diagnosis per 2010 ACR Criteria; added Appendix C to define MTX failure. NOMID: Added weight based dosing limit.	08.17	08.17
2Q 2018 annual review: added HIM; removed TB testing requirement from all indications; references reviewed and updated.	02.27.18	05.18

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan

retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note:**

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

**For Health Insurance Marketplace members**, when applicable, this policy applies only when the prescribed agent is on your health plan approved formulary. Request for non-formulary drugs must be reviewed using the formulary exception policy.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.