

Clinical Policy: Sacubitril/Valsartan (Entresto)

Reference Number: CP.PMN.67

Effective Date: 11.01.15

Last Review Date: 02.19

Line of Business: Commercial, HIM, Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Sacubitril/valsartan (Entresto[®]) is a combination of sacubitril, a neprilysin inhibitor, and valsartan, an angiotensin II receptor blocker (ARB).

FDA Approved Indication(s)

Entresto is indicated to reduce the risk of cardiovascular death and hospitalization for heart failure in patients with chronic heart failure (New York Heart Association [NYHA] Class II-IV) and reduced ejection fraction.

Entresto is usually administered in conjunction with other heart failure therapies, in place of an angiotensin-converting enzyme (ACE) inhibitor or other ARB.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Entresto is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Heart Failure (must meet all):

1. Diagnosis of chronic heart failure of NYHA Class II, III, or IV;
2. Age \geq 18 years;
3. Left ventricular ejection fraction (LVEF) is \leq 35%;
4. At the time of request, member has none of the following contraindications:
 - a. Concomitant use with ACE inhibitors;
 - b. If member has a diagnosis of diabetes, concomitant use with aliskiren;
5. Dose does not exceed sacubitril 194 mg/valsartan 206 mg (2 tablets) per day.

Approval duration:

HIM – 12 months

Commercial/Medicaid – Length of Benefit

B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Heart Failure (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Entresto for heart failure and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed sacubitril 194 mg/valsartan 206 mg (2 tablets) per day.

Approval duration:

HIM – 12 months

Commercial/Medicaid – Length of Benefit

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

Approval duration: Duration of request or 12 months (whichever is less); or

2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- ### A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

ACE: angiotensin-converting enzyme

ARB: angiotensin II receptor blocker

FDA: Food and Drug Administration

NYHA: New York Heart Association

LVEF: left ventricular ejection fraction

Appendix B: Therapeutic Alternatives

Not applicable

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s):
 - Hypersensitivity to any component
 - History of angioedema related to previous ACE inhibitor or ARB therapy
 - Concomitant use of Entresto with an ACE inhibitor is contraindicated because of the increased risk of angioedema.
 - Concomitant use of Entresto and ARB should be avoided since Entresto contains an ARB.

- Boxed warning(s): When pregnancy is detected, discontinue ENTRESTO as soon as possible. Drugs that act directly on the renin-angiotensin system can cause injury and death to the developing fetus.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Chronic heart failure	<p>The recommended starting dose is 49/51 mg (sacubitril/valsartan) PO BID. Double the dose after 2 to 4 weeks to the target maintenance dose of 97/103 mg (sacubitril/valsartan) BID, as tolerated by the patient.</p> <p>Therapy may be initiated at 24/26 mg (sacubitril/valsartan) PO BID for:</p> <ul style="list-style-type: none"> • patients not currently taking an ACE inhibitor or an ARB or previously taking a low dose of these agents • patients with severe renal impairment • patients with moderate hepatic impairment <p>Double the dose every 2 to 4 weeks to the target maintenance dose of 97/103 mg (sacubitril/valsartan) BID, as tolerated by the patient.</p>	Sacubitril 194 mg/valsartan 206 mg per day

VI. Product Availability

Film-coated tablets (sacubitril/valsartan): 24 mg/26 mg, 49 mg /51 mg, 97 mg/103 mg

VII. References

1. Entresto Prescribing Information. East Hanover, NJ: Novartis Pharmaceuticals Corporation; November 2017. Available at: <https://www.pharma.us.novartis.com/sites/www.pharma.us.novartis.com/files/entresto.pdf>. Accessed October 30, 2018.
2. Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Colvin MM, Drazner MH, Filippatos GS, Fonarow GC, Givertz MM, Hollenberg SM, Lindenfeld J, Masoudi FA, McBride PE, Peterson PN, Stevenson LW, Westlake C. 2017 ACC/AHA/HFSA focused update of the 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology /American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *J Am Coll Cardiol*. 2017.
3. Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Colvin MM, Drazner MH, Filippatos G, Fonarow GC, Givertz MM, Hollenberg SM, Lindenfeld J, Masoudi FA, McBride PE, Peterson PN, Stevenson LW, Westlake C. 2016 ACC/AHA/HFSA focused update on new pharmacological therapy for heart failure: an update of the 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *Circulation*. 2016;134: 000-000.
4. Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Drazner MH, Fonarow GC, Geraci SA, Horwich T, Januzzi JL, Johnson MR, Kasper EK, Levy WC, Masoudi FA, McBride PE,

- McMurray JJ, Mitchell JE, Peterson PN, Riegel B, Sam F, Stevenson LW, Tang WH, Tsai EJ, Wilkoff BL; American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation*. 2013 Oct 15;128(16):e240-327.
6. McMurray JJ, Packer M, Desai AS, et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med*. 2014;371:993-1004.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
New guideline created	09.15	11.15
Converted to new integrated template. Removed age requirement since not referenced in indications section in PI. Modified specific max quantity limit to FDA max recommended dose and health plan approved QL statement. Increased initial approval duration to 12 months. Removed “For renewal request below the target dose of sacubitril/valsartan 97/103mg twice daily, provider must provide a clinical rationale for continued treatment at a sub-therapeutic dose” as dose is dependent on patient tolerability per PI and provider’s clinical judgment. Updated continuation criteria to include continuity of care Updated references to reflect current literature search	08.16	11.16
Converted to new template. Added age restriction and contraindications related to DDI per PI/safety approach. Modified max dose requirement to include specific quantity limit. Updated references.	08.07.17	11.17
1Q18 annual review. Policies combined for Centene Medicaid, Marketplace and Commercial lines of business; No significant change from previous corporate approved policy; Commercial: removed prescriber requirement; added age restriction as safety and effectiveness in pediatric patients have not been established; modified LVEF from < 40% to ≤ 35% per PARADIGM-HF clinical trial; added contraindications related to DDI per PI; updated re-auth to allow COC for heart failure. Added requirement for positive response to therapy; Marketplace and Medicaid: removed prescriber requirement; added age restriction and contraindications related to DDI per PI (Marketplace only); removed “previously tolerated an ACEI or ARB at therapeutic doses for ≥ 30 days” since specialist is involved in care; References reviewed and updated.	11.01.17	02.18
Medicaid: approval duration changed from 12 months to length of benefit	03.04.18	05.18
1Q 2019 annual review: no significant changes; references reviewed and updated.	10.30.18	02.19

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members

and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

For Health Insurance Marketplace members, when applicable, this policy applies only when the prescribed agent is on your health plan approved formulary. Request for non-formulary drugs must be reviewed using the formulary exception policy; HIM.PA.103.

©2015 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.