

## Clinical Policy: Dipeptidyl Peptidase-4 (DPP-4) Inhibitors

Reference Number: HIM.PA.58

Effective Date: 03.01.18

Last Review Date: 02.23

Line of Business: HIM

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### Description

The following agents contain a dipeptidyl peptidase-4 (DPP-4) inhibitor and require prior authorization\*: alogliptin (Nesina<sup>®</sup>), alogliptin/metformin (Kazano<sup>®</sup>), alogliptin/pioglitazone (Oseni<sup>®</sup>), linagliptin (Tradjenta<sup>®</sup>), linagliptin/metformin (Jentadueto<sup>®</sup>, Jentadueto<sup>®</sup> XR), saxagliptin (Onglyza<sup>®</sup>), and saxagliptin/metformin (Kombiglyze XR<sup>®</sup>).

*\*If request is for a combination DPP-4 inhibitor and sodium glucose co-transporter 2 (SGLT2) inhibitor (e.g., linagliptin/empagliflozin [Glyxambi<sup>®</sup>], linagliptin/empagliflozin /metformin [Trijardy<sup>™</sup> XR], saxagliptin/dapagliflozin [Qtern<sup>®</sup>], sitagliptin/ertugliflozin [Steglujan<sup>™</sup>]), refer to HIM.PA.91 SGLT2 Inhibitors.*

### FDA Approved Indication(s)

DPP-4 inhibitors are indicated as adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus.

Limitation(s) of use:

- DPP-4 inhibitors should not be used in patients with type 1 diabetes.
- Onglyza and Kombiglyze should not be used for the treatment of diabetic ketoacidosis.
- Tradjenta, Jentadueto, and Jentadueto XR have not been studied in patients with a history of pancreatitis.

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that DPP-4 inhibitors are **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Type 2 Diabetes Mellitus (must meet all):

1. Diagnosis of type 2 diabetes mellitus;
2. Age  $\geq$  18 years;
3. Member meets one of the following (a or b):
  - a. Failure of  $\geq$  3 consecutive months of metformin, unless contraindicated or clinically significant adverse effects are experienced;
  - b. For antidiabetic medication-naïve members, requested agent is approvable if intended for concurrent use with metformin due to HbA1c  $\geq$  8.5% (drawn within the past 3 months);

4. Failure of  $\geq 3$  consecutive months of a sitagliptin-containing product (e.g., sitagliptin [Januvia<sup>®</sup>], sitagliptin/metformin [Janumet<sup>®</sup>, Janumet<sup>®</sup> XR]), unless clinically significant adverse effects are experienced or all are contraindicated;
5. Dose does not exceed the FDA approved maximum recommended dose (*see Section V*).

**Approval duration: 12 months**

**B. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: HIM.PA.33 for health insurance marketplace; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: HIM.PA.103 for health insurance marketplace; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: HIM.PA.154 for health insurance marketplace.

**II. Continued Therapy**

**A. Type 2 Diabetes Mellitus (must meet all):**

1. Member meets one of the following (a or b):
  - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
  - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed the FDA approved maximum recommended dose (*see Section V*).

**Approval duration: 12 months**

**B. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: HIM.PA.33 for health insurance marketplace; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: HIM.PA.103 for health insurance marketplace; or

2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: HIM.PA.154 for health insurance marketplace.

**III. Diagnoses/Indications for which coverage is NOT authorized:** Not applicable

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – HIM.PA.154 for health insurance marketplace or evidence of coverage documents.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

AACE: American Association of Clinical Endocrinologists

ACE: American College of Endocrinology

ADA: American Diabetes Association

ASCVD: atherosclerotic cardiovascular disease

DPP-4: dipeptidyl peptidase-4

FDA: Food and Drug Administration

GLP-1: glucagon-like peptide-1

HbA1c: glycated hemoglobin

HF: heart failure

SGLT2: sodium-glucose co-transporter 2

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

<b>Drug Name</b>	<b>Dosing Regimen</b>	<b>Dose Limit/ Maximum Dose</b>
metformin (Fortamet <sup>®</sup> , Glucophage <sup>®</sup> , Glucophage <sup>®</sup> XR, Glumetza <sup>®</sup> )	Regular-release (Glucophage): 500 mg PO BID or 850 mg PO QD; increase as needed in increments of 500 mg/week or 850 mg every 2 weeks  Extended-release: <ul style="list-style-type: none"> <li>• Fortamet, Glumetza: 1,000 mg PO QD; increase as needed in increments of 500 mg/week</li> <li>• Glucophage XR: 500 mg PO QD; increase as needed in increments of 500 mg/week</li> </ul>	Regular-release: 2,550 mg/day  Extended-release: 2,000 mg/day
Januvia (sitagliptin)	100 mg PO QD	100 mg/day
Janumet (sitagliptin/metformin)	Individualized dose PO BID	100/2,000 mg/day
Janumet XR (sitagliptin/metformin)	Individualized dose PO QD	100/2,000 mg/day

*Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.*

*Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s):
  - History of serious hypersensitivity reaction to the requested drug product
  - Severe renal impairment (*metformin-containing products*)
  - Acute or chronic metabolic acidosis, including diabetic ketoacidosis (*metformin-containing products only*)
  - NYHA Class III or IV heart failure (*Oseni only*)
- Boxed warning(s): lactic acidosis (*metformin-containing products only*), congestive heart failure (*Oseni only*)

*Appendix D: General Information*

- Per the American Diabetes Association (ADA) and American Association of Clinical Endocrinologists and American College of Endocrinology (AAACE/ACE) guidelines:
  - Metformin is recommended for all patients with type 2 diabetes. It is effective and safe, is inexpensive, and may reduce risk of cardiovascular events and death. Monotherapy is recommended for most patients; however:
    - Starting with dual therapy (i.e., metformin plus another agent, such as a sulfonylurea, thiazolidinedione, DPP-4 inhibitor, SGLT2 inhibitor, glucagon-like peptide 1 [GLP-1] receptor agonist, or basal insulin) may be considered for patients with baseline HbA1c  $\geq 1.5\%$  above their target. According to the ADA, a reasonable HbA1c target for many non-pregnant adults is  $< 7\%$  ( $\leq 6.5\%$  per the AAACE/ACE).
    - Starting with combination therapy with insulin may be considered for patients with baseline HbA1c  $> 10\%$  or if symptoms of hyperglycemia are present.
  - If the target HbA1c is not achieved after approximately 3 months of monotherapy, dual therapy should be initiated. If dual therapy is inadequate after 3 months, triple therapy should be initiated. Finally, if triple therapy fails to bring a patient to goal, combination therapy with insulin should be initiated. Each non-insulin agent added to initial therapy can lower HbA1c by 0.7-1%.

**V. Dosage and Administration**

Drug Name	Dosing Regimen	Maximum Dose
Jentadueto (linagliptin/metformin)	Individualized dose PO BID	5/2,000 mg/day
Jentadueto XR (linagliptin/metformin)	Individualized dose PO QD	5/2,000 mg/day
Kazano (alogliptin/metformin)	Individualized dose PO BID	25/2,000 mg/day
Kombiglyze XR (saxagliptin/metformin)	Individualized dose PO QD	5/2,000 mg/day
Nesina (alogliptin)	25 mg PO QD	25 mg/day
Onglyza (saxagliptin)	2.5 or 5 mg PO QD	5 mg/day
Oseni (alogliptin/pioglitazone)	Individualized dose PO QD	25/45 mg/day
Tradjenta (linagliptin)	5 mg PO QD	5 mg/day

**VI. Product Availability**

Drug Name	Availability
Jentadueto (linagliptin/metformin)	Tablets: 2.5/500 mg, 2.5/850 mg, 2.5/1,000 mg
Jentadueto XR (linagliptin/metformin)	Tablets: 5/1,000 mg, 2.5/1,000 mg

Drug Name	Availability
Kazano (alogliptin/metformin)	Tablets: 12.5/500 mg, 12.5/1,000 mg
Kombiglyze XR (saxagliptin/metformin)	Tablets: 5/500 mg, 5/1,000 mg, 2.5/1,000 mg
Nesina (alogliptin)	Tablets: 6.25 mg, 12.5 mg, 25 mg
Onglyza (saxagliptin)	Tablets: 2.5 mg, 5 mg
Oseni (alogliptin/pioglitazone)	Tablets: 12.5/15 mg, 12.5/30 mg, 12.5/45 mg, 25/15 mg, 25/30 mg, 25/45 mg
Tradjenta (linagliptin)	Tablets: 5 mg

## VII. References

- American Diabetes Association. Standards of medical care in diabetes—2022. *Diabetes Care*. 2022; 45(suppl 1): S1-S264. Updated May 2022. Accessed October 26, 2022.
- Blonde L, Umpierrez GE, Reddy SS, et al. American Association of Clinical Endocrinology clinical practice guideline: Developing a diabetes mellitus comprehensive care plan – 2022 update. *Endocrine Practice*. 2022; 28(10): 923-1049.
- Jentaduetto Prescribing Information Ridgefield, CT: Boehringer Ingelheim Pharmaceuticals, Inc.; April 2022. Available at: [www.jentaduetto.com](http://www.jentaduetto.com). Accessed October 27, 2022.
- Jentaduetto XR Prescribing Information. Ridgefield, CT: Boehringer Ingelheim Pharmaceuticals, Inc.; October 2021. Available at: [www.jentaduettoxr.com](http://www.jentaduettoxr.com). Accessed October 27, 2022.
- Kombiglyze XR Prescribing Information. Wilmington, DE: AstraZeneca Pharmaceuticals LP; October 2019. Available at: [www.kombiglyzextr.com](http://www.kombiglyzextr.com). Accessed October 27, 2022.
- Onglyza Prescribing Information. Wilmington, DE: AstraZeneca Pharmaceuticals LP; October 2019. Available at: [www.onglyza.com](http://www.onglyza.com). Accessed October 27, 2022.
- Tradjenta Prescribing Information. Ridgefield, CT: Boehringer Ingelheim Pharmaceuticals, Inc.; April 2022. Available at: [www.tradjenta.com](http://www.tradjenta.com). Accessed October 27, 2022.
- Nesina Prescribing Information. Deerfield, IL: Takeda Pharmaceuticals America, Inc.; March 2022. Available at: [www.nesinafamily.com](http://www.nesinafamily.com). Accessed October 27, 2022.
- Oseni Prescribing Information. Deerfield, IL: Takeda Pharmaceuticals America, Inc.; March 2022. Available at: [www.nesinafamily.com](http://www.nesinafamily.com). Accessed October 27, 2022.
- Kazano Prescribing Information. Deerfield, IL: Takeda Pharmaceuticals America, Inc.; March 2022. Available at: [www.nesinafamily.com](http://www.nesinafamily.com). Accessed October 27, 2022.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
1Q 2019 annual review: modified minimum A1c related for concurrent use of metformin from 9% to 8.5% based on 2019 ADA guidelines; references reviewed and updated.	11.01.18	02.19
Added requirement for trial of Steglatro or Segluromet prior to Glyxambi to align with criteria for Glyxambi in the SGLT2 clinical policy; members requesting other non-preferred DPP-4 inhibitors are still required to try/fail the preferred DPP-4 inhibitors Tradjenta and Januvia.	04.22.19	

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Per SDC and prior clinical guidance added Onglyza to criteria requiring redirection to the preferred DPP-4 inhibitors (sitagliptin or linagliptin-containing products, which include the addition of Janumet/XR and Jentadueto/XR); applied similar redirection to Nesina.	10.23.19	
1Q 2020 annual review: no significant changes; added Trijardy XR with re-direction to Steglatro or Segluromet per SDC; references reviewed and updated.	10.29.19	02.20
Allowed bypass of Steglatro/Segluromet for patients with established cardiovascular disease or diabetic nephropathy requesting Glyxambi/Trijardy XR per previously approved clinical guidance and SDC clarification.	04.01.20	
Per September SDC and prior clinical guidance for 2021, added Steglujan and applied revised Glyxambi and Trijardy XR redirection to require an empagliflozin, ertugliflozin, or sitagliptin-containing product; revised Nesina and Onglyza redirection to require sitagliptin-containing product only (removed redirection to linagliptin-containing product) and applied similar redirection to Tradjenta, Jentadueto, or Jentadueto XR which were added to the policy; added Oseni, Kazano and Kombiglyze XR to policy.	09.08.20	
Per December SDC and prior clinical guidance, added specific redirection to Glyxambi or Trijardy XR for Steglujan, removed Glyxambi and Trijardy XR from policy as prior authorization is not required.	12.15.20	
1Q 2021 annual review: removed criteria for combination DPP4/SGLT2 products and directed requests to the SGLT2 policy instead; references reviewed and updated.	10.27.20	02.21
1Q 2022 annual review: no significant changes; references reviewed and updated.	09.16.21	02.22
Template changes applied to other diagnoses/indications and continued therapy section.	10.11.22	
1Q 2023 annual review: no significant changes; references reviewed and updated.	10.26.22	02.23

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical

practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.